


Study examines immune responses in patients with kidney failure after receiving different COVID-19 vaccines

November 23 2021



Credit: CC0 Public Domain

Death rates from COVID-19 are especially high in individuals with kidney failure who are on dialysis, making SARS-CoV-2 vaccination in this population a high priority. A recent study compared the immune

responses of patients with and without kidney failure following immunization with different COVID-19 vaccines. The findings, which are published in *JASN*, may help improve vaccination strategies in vulnerable patients.

Individuals on dialysis typically have an impaired response to vaccination. To better understand the predictors and dynamics of their antibody and cellular immune responses to different SARS-CoV-2 vaccines, a team led by An S. De Vriese, MD, Ph.D. (AZ Sint-Jan Brugge, in Belgium) prospectively evaluated responses at 4 or 5 weeks and again at 8 or 9 weeks after immunization with the Pfizer-BioNTech (BNT162b2) and Moderna (mRNA-1273) mRNA vaccines in 543 patients on hemodialysis and 75 individuals with normal [kidney](#) function.

In the multicenter study, the researchers found an incomplete and delayed antibody response and a blunted cellular immune response to vaccination in patients on hemodialysis. However, responses were substantially stronger with the mRNA-1273 vaccine than with the BNT162b2 vaccine (in both individuals with [kidney failure](#) and those with normal kidney function). The researchers believe that this may be due to the higher mRNA dose in the mRNA-1273 vaccine than in the BNT162b2 vaccine. "These differences in immune [response](#) may translate into differences in vaccine effectiveness in vulnerable populations in the long-term, when protective immunity is waning and in the battle against the Delta variant," said Dr. De Vriese.

In addition, patients on hemodialysis who had a prior COVID-19 infection, did not take immunosuppressive drugs, had higher serum albumin levels and lymphocyte counts, had previously responded to hepatitis B vaccination, and were on dialysis for only a short amount of time had higher antibody and cellular immune responses.

"We believe that a high-dose vaccine may be a valid strategy to improve SARS-CoV-2 vaccine effectiveness in hemodialysis patients. The most vulnerable patients—those who are using [immunosuppressive drugs](#), have a low serum albumin, a low lymphocyte count, are hepatitis B vaccine non-responders, or have a high dialysis vintage—may be good candidates for a third [vaccine](#) dose," said Dr. De Vriese.

Provided by American Society of Nephrology

Citation: Study examines immune responses in patients with kidney failure after receiving different COVID-19 vaccines (2021, November 23) retrieved 10 May 2024 from <https://medicalxpress.com/news/2021-11-immune-responses-patients-kidney-failure.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--