

# More kids to be tested for lead poisoning, but eliminating sources is key

November 8 2021, by Gabriel Filippelli

---



Credit: Unsplash/CC0 Public Domain

The U.S. Centers for Disease Control and Prevention has [updated its blood lead reference value](#)—the level at which children ages 1–5 are considered to have high exposure to lead. Since 2012, this threshold had

been set at 5 micrograms of lead per deciliter of blood; children at or above this level represented the top 2.5% with the highest blood lead levels in the nation. Now, in response to recent federal health surveys, the CDC has updated that number to 3.5 micrograms per deciliter. Environmental scientist [Gabriel Filippelli](#), who has studied urban lead poisoning in children, explains what this shift means for public health.

## **Will this change affect how doctors detect and treat childhood lead poisoning?**

The Centers for Disease Control periodically reviews national data on blood lead levels in [children](#). This new lower value is the [average blood lead level exceeded by 2.5% of children tested](#).

Many clinics have an on-site screening device that uses electrochemical detection to quickly test a small amount of blood from a fingertip prick. If children test positive, doctors refer them to have a larger blood sample drawn from a vein and analyzed in a diagnostic laboratory. The [clinical test](#) is fast, cheap and relatively painless, but the venous blood draw is the gold standard for diagnosing lead poisoning.

On-site clinical devices typically can detect lead at concentrations as low as 3.2 micrograms per deciliter, so the new CDC guidance means that nearly all children who show positive results at the screening level will be referred for follow-up testing. That's much more protective from a public health perspective.

However, it will [roughly double](#) the number of children who are classified as at highest risk for lead poisoning. Formerly, children had to have at least 5 micrograms per deciliter of lead in their blood to fall into that group; now it will include thousands more children with slightly lower blood lead levels.

Larger numbers of children means that many states will have trouble affording testing and follow-up care—which can involve [dietary changes and medications](#), as well as removing lead exposure sources—unless Congress increases federal support for programs to prevent and treat lead poisoning.

This [#BabySafetyMonth](#), remind your doctor to test your child for lead poisoning. Health care providers are required to test all 1- and 2-year-old children for lead. Children should also be assessed for lead exposure every year until they are 6 years old: <https://t.co/CQhB4o8fJX> [pic.twitter.com/fqcMnySPHF](https://t.co/fqcMnySPHF)

— nychealthy (@nycHealthy) [September 13, 2021](#)

## How are children commonly exposed to lead?

The most pervasive source, especially in cities, is soil and dust generated from soil. Thanks to many years of emissions from degraded lead-based paint, leaded gasoline and industrial sources, typical urban soils have lead concentrations that [range from benign to toxic](#). Children are exposed when they touch or play in contaminated dirt or inhale the dust.

The U.S. Environmental Protection Agency's limit for lead in soils in public play areas is [400 parts per million](#). That's significantly higher than typical background levels, which are roughly 20 to 50 parts per million.

Some U.S. states, [such as California](#), have much lower limits. In my experience, it's not unusual to find urban soils with much higher levels, particularly near the exterior walls of buildings where lead may accumulate from degraded paints or [dust buildup](#).

The most lead-contaminated neighborhoods in cities are often the poorest and home to the [highest percentage of nonwhite children](#). This is

a legacy of [racist housing practices](#) that concentrated people of color in less desirable neighborhoods. Residents in these zones can have significantly higher rates of elevated blood lead levels than people in wealthier neighborhoods.

Lead-based paint is also a major exposure risk, particularly in poorly maintained buildings. Lead paint tastes sweet, so children sometimes chew on paint chips or painted wood.

Lead water pipes are a third source, although less common than paint or soil. Many cities and towns across the U.S. have lead service lines that deliver water to homes. If their water is treated properly, a protective plaque forms on the inside of water pipes and seals their lead content away from the water.

But some cities, including [Washington, D.C.](#), [Newark](#), New Jersey, and [Flint, Michigan](#), have changed their water sources or treatment processes in ways that stripped out the protective plaque and carried lead to household taps. These water crises disproportionately affected communities of color.

## **How does lead exposure at these levels affect children's health?**

Historically, public health interventions focused on acutely poisoned children who exhibited clear neurocognitive issues such as attention deficit, memory lapses, agitation and even tremors. As lead was slowly removed from most home uses in the mid-20th century and the U.S. population's blood lead levels decreased, these obvious clinical presentations for lead poisoning declined.

What we see now are more subtle neurocognitive deficits, which

scientists and medical experts measure through neurological and behavioral testing. A child who is diagnosed as having high blood lead levels today may perform poorly on standardized exams, behave disruptively in the classroom or at home or have trouble retaining information. Follow-up research in Flint shows that many infants and toddlers who were exposed to lead in water there in 2015 [are struggling now that they are in school](#).

These types of tests show that blood lead levels even lower than the new standard [still affect performance](#). This research is the basis for statements from scholars and the CDC that [there is no safe blood lead level in children](#).

## **What's the trend for childhood lead poisoning in the US?**

It has been falling since most major environmental sources of lead, such as leaded gasoline, lead-based paints and industrial emissions, were eliminated starting in the 1970s. Recent analyses show that the median blood lead level for all U.S. children between ages 1 and 5 is about 0.7 micrograms per deciliter today, compared with [15 micrograms per deciliter in the late 1970s](#).

But Black children and children living in poverty have average blood lead levels that are [13% higher than this national average](#), which means that many of them are at risk.

For example, in a 2019 study, I worked with colleagues at Notre Dame to analyze blood lead levels of over 18,000 children in St. Joseph County, Indiana, which includes the town of South Bend. In some neighborhoods, over 30% of children had [blood](#) lead levels higher than 5 micrograms per deciliter, and over 65% of the census tracts had average

[blood lead levels over that safety limit.](#)

We also found that there was no systematic, risk-informed approach to testing. In areas that had the highest potential risks based on poverty levels, less than 6% of eligible children had lead test results reported to the county health department—the same rate as in other, wealthier census tracts. Without more screening, and more work to eliminate lead exposure in the communities most at risk, this problem won't be solved for a long time.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: More kids to be tested for lead poisoning, but eliminating sources is key (2021, November 8) retrieved 10 April 2024 from <https://medicalxpress.com/news/2021-11-kids-poisoning-sources-key.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--