


# Could EKGs help doctors use AI to detect pulmonary embolisms?

December 21 2021

---



In a pilot study, researchers at the Icahn School of Medicine at Mount Sinai showed that AI algorithms that rely on electrocardiogram and electronica health

records may one day help doctors screen for pulmonary embolisms. Credit: Glicksberg and Nadkarni labs, Mount Sinai, N.Y., N.Y.

Pulmonary embolisms are dangerous, lung-clogging blot clots. In a pilot study, scientists at the Icahn School of Medicine at Mount Sinai have shown for the first time that artificial intelligence (AI) algorithms can detect signs of these clots in electrocardiograms (EKGs), a finding that may one day help doctors with screening.

The results published in the *European Heart Journal-Digital Health* suggested that new machine learning algorithms, which are designed to exploit a combination of EKG and electronic [health](#) record (EHR) data, may be more effective than currently used screening tests at determining whether moderate- to [high-risk patients](#) actually have [pulmonary embolisms](#).

The study was led by Sulaiman S. Somani, MD, a former medical student in the lab of Benjamin S. Glicksberg, Ph.D., Assistant Professor of Genetics and Genomic Sciences and a member of the Hasso Plattner Institute for Digital Health at Mount Sinai.

Pulmonary embolisms happen when deep vein blood clots, usually formed in the legs or arms, break away and clog lung arteries. These clots can be lethal or cause long-term lung damage. Although some patients may experience shortness of breath or chest pain, these symptoms may also signal other problems that have nothing to do with blood clots, making it difficult for doctors to properly diagnose and treat cases. Moreover, current official diagnoses rely on computed tomography pulmonary angiograms (CTPAs), which are time-consuming chest scans that can only be performed at select hospitals and require patients to be exposed to potentially dangerous levels of radiation.

To make diagnoses easier and more accessible, researchers have spent more than 20 years developing advanced computer programs, or algorithms, designed to help doctors determine whether at-risk patients are actually experiencing pulmonary embolisms. The results have been mixed. For example, algorithms that used EHRs have produced a wide range of success rates for accurately detecting clots and can be labor-intensive. Meanwhile, the more accurate ones depend heavily on data from the CTPAs.

In this study the researchers found that fusing algorithms that rely on EKG and EHR data may be an effective alternative, because EKGs are widely available and relatively easy to administer.

The researchers created and tested out various algorithms on data from 21,183 Mount Sinai Health System patients who showed moderate to highly suspicious signs of having pulmonary embolisms. While some algorithms were designed to use EKG data to screen for pulmonary embolisms, others were designed to use EHR data. In each situation, the [algorithm](#) learned to identify a pulmonary embolism case by comparing either EKG or EHR data with corresponding results from CTPAs. Finally, a third, fusion algorithm was created by combining the best-performing EKG algorithm with the best-performing EHR one.

The results showed that the fusion model not only outperformed its parent algorithms but was also better at identifying specific pulmonary embolism cases than the Wells' Criteria Revised Geneva Score and three other currently used screening tests. The researchers estimated that the fusion model was anywhere from 15 to 30 percent more effective at accurately screening acute embolism cases, and the model performed best at predicting the most severe cases. Furthermore, the fusion model's accuracy remained consistent regardless of whether race or sex was tested as a factor, suggesting it may be useful for screening a variety of patients.

According to the authors, these results support the theory that EKG data may be effectively incorporated into new pulmonary embolism screening algorithms. They plan to further develop and test these algorithms out for potential utility in the clinic.

**More information:** Sulaiman S Somani et al, Development of a machine learning model using electrocardiogram signals to improve acute pulmonary embolism screening, *European Heart Journal-Digital Health* (2021). [DOI: 10.1093/ehjdh/ztab101](https://doi.org/10.1093/ehjdh/ztab101)

Provided by The Mount Sinai Hospital

Citation: Could EKGs help doctors use AI to detect pulmonary embolisms? (2021, December 21) retrieved 2 May 2024 from <https://medicalxpress.com/news/2021-12-ekgs-doctors-ai-pulmonary-embolisms.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--