


People with IBD have more microplastics in their feces, study says

December 22 2021



Higher numbers of microplastics of various shapes, such as sheets (left) and fibers (right) were found in the feces of people with IBD than in healthy controls. Credit: Adapted from *Environmental Science & Technology* 2021, DOI: 10.1021/acs.est.1c03924

Microplastics—tiny pieces of plastic less than 5 mm in length — are everywhere, from bottled water to food to air. According to recent estimates, people consume tens of thousands of these particles each year, with unknown health consequences. Now, researchers reporting in *ACS' Environmental Science & Technology* found that people with

inflammatory bowel disease (IBD) have more microplastics in their feces than healthy controls, suggesting that the fragments could be related to the disease process.

The prevalence of IBD, which includes Crohn's disease and ulcerative colitis, is rising globally. Characterized by chronic inflammation of the digestive tract, IBD can be triggered or made worse by diet and environmental factors. Microplastics can cause intestinal inflammation, gut microbiome disturbances and other problems in animal models, so Faming Zhang, Yan Zhang and colleagues wondered if they could also contribute to IBD. As a first step toward finding out, the researchers wanted to compare the levels of microplastics in feces from healthy subjects and people with different severities of IBD.

The team obtained [fecal samples](#) from 50 healthy people and 52 people with IBD from different geographic regions of China. Analysis of the samples showed that feces from IBD patients contained about 1.5 times more [microplastic](#) particles per gram than those from healthy subjects. The microplastics had similar shapes (mostly sheets and fibers) in the two groups, but the IBD feces had more small (less than 50 μm) particles. The two most common types of plastic in both groups were polyethylene terephthalate (PET; used in bottles and [food](#) containers) and polyamide (PA; found in food packaging and textiles). People with more severe IBD symptoms tended to have higher levels of fecal microplastics. Through a questionnaire, the researchers found that people in both groups who drank bottled water, ate takeaway food and were often exposed to dust had more microplastics in their feces.

These results suggest that people with IBD may be exposed to more microplastics in their gastrointestinal tract. However, it's still unclear whether this exposure could cause or contribute to IBD, or whether people with IBD accumulate more fecal microplastics as a result of their disease, the researcher say.

More information: Analysis of Microplastics in Human Feces Reveals a Correlation between Fecal Microplastics and Inflammatory Bowel Disease Status, *Environmental Science & Technology* (2021). [DOI: 10.1021/acs.est.1c03924](https://doi.org/10.1021/acs.est.1c03924)

Provided by American Chemical Society

Citation: People with IBD have more microplastics in their feces, study says (2021, December 22) retrieved 18 April 2024 from <https://medicalxpress.com/news/2021-12-people-ibd-microplastics-feces.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.