

Scientists develop an RNA-based breath test to detect COVID-19

December 13 2021



Bubbler kits were processed in a negative pressure bench top box to prevent contamination of amplicon particles in the laboratory. Credit: William G. Fairbrother

In a new study in *The Journal of Molecular Diagnostics*, investigators report on the design and testing of a breathalyzer, known as the Bubbler,

that relies on viral RNA detection to diagnose SARS-CoV-2. Its name is derived from the bubbling sound that occurs when the patient exhales into the device.


The Bubbler not only reverse transcribes RNA from airborne virus particles into DNA to be tested via PCR but can also barcode that DNA, allowing samples to be linked directly to the patient they have come from and be used for sequencing. It can be used for simultaneous batches of pooled samples and provides additional information such as [viral load](#) and strain identity and eliminates the need for stabilizing a sample, potentially allowing the assay to be performed at home.

"Involvement of the lower [respiratory tract](#) is often a precursor to severe COVID-19, so there is an argument for a more direct sampling focused on exhaled breath," explained lead investigator William G. Fairbrother, Ph.D., professor in the department of molecular biology, [cell biology](#) and biochemistry at Brown University in Providence, RI, U.S.

Virus detection by the Bubbler is similar to a hospital-swab PCR test; however, it is a better measure of risk of contagion as it detects airborne viral particles. Swab tests can return a positive result for months after infection as they detect viral RNA fragments in cells that persist in previously infected cells. The Bubbler can also be adapted for environmental sampling in hospitals, transportation hubs, and closed environments like offices, ships, and planes, the investigators report.

Seventy patients treated in the Emergency Department of Rhode Island Hospital between May 2020 and January 2021 were screened. The study tested samples from three points in the respiratory tract. Tongue scrapes from the mouth (saliva/tongue scrapes) and from 15 seconds of exhaled breath collected in the Bubbler were compared to those from a conventional nasopharyngeal swab PCR test. The Bubbler is a [glass tube](#) with a glass pipette through which patients can exhale. The tube is filled

with a reverse transcription reaction mixture and cold mineral oil.



Breathalyzers were bathed in UV light to maintain sterile field prior to construction of kits prepared for the clinical trial at Rhode Island Hospital.
Credit: William G. Fairbrother

The study determined that SARS-CoV-2 can be readily detected in the breath and is more predictive of lower respiratory tract involvement. Viral RNA is more enriched in the breath relative to oral samples, while oral samples include cells involved with SARS-CoV-2 replication that breath samples do not. This suggests the viral signal detected in the Bubbler comes from active viral particles.

"The Bubbler is more likely to be a better indicator of current infection than nasopharyngeal swabs," said Dr. Fairbrother. "Another advantage is the barcoding, which enables high-throughput RNA virus testing at a fraction of the cost of conventional testing. The barcode returns a viral sequence that also supports strain identification, which may prove useful as more information is learned about transmissibility and possible strain-specific treatment decisions."

The investigators also demonstrated how the Bubbler might be adapted to detect virus in airborne samples. To model the movement of droplets exhaled in human breath, three unique nucleic acid samples were added to three personal humidifiers at different locations at varied distances from the Bubbler in a room with high airflow and a room with low airflow. Although a detailed exploration of this application was beyond the scope of the study, the results demonstrate the potential to use aerosolized nucleic acids to quantitatively map airflow in indoor spaces, and to detect SARS-CoV-2 in the air.

"Such technology could be useful in restoring service to industries such as hotels, cruise ships, and casinos," Dr. Fairbrother observed. "There is also an epidemiological benefit to routine testing of air at early warning sites such as transportation hubs and hospital emergency departments."

Tests for COVID-19 usually use samples collected from the upper

respiratory tract by saliva or nasopharyngeal swab. Positive samples contain active virus, but viral load in the upper respiratory tract is not correlated with symptoms in the lower respiratory tract, such as pneumonia.

More information: Chaorui Duan et al, Efficient Detection of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) from Exhaled Breath, *The Journal of Molecular Diagnostics* (2021). [DOI: 10.1016/j.jmoldx.2021.09.005](https://doi.org/10.1016/j.jmoldx.2021.09.005)

Provided by Elsevier

Citation: Scientists develop an RNA-based breath test to detect COVID-19 (2021, December 13) retrieved 29 June 2024 from <https://medicalxpress.com/news/2021-12-scientists-rna-based-covid-.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.