

# How can body weight affect the mortality risk of excessive drinkers?

January 24 2022

---



Credit: Unsplash/CC0 Public Domain

While research has long shown a higher risk of death linked to alcoholism for people with overweight, a new study published in the journal *Drug and Alcohol Dependence* has found that people with

underweight who drink excessively may be at an even higher risk of dying from heart disease, cancer and other causes.

The study was based on data from the National Health Interview Survey (NHIS), which has a nationally representative sample of more than 200,000 U.S. adults aged 35-85, interviewed between Jan. 1, 2001, and Dec. 31, 2011. The researchers analyzed data on [mortality risk](#) among drinkers and non-drinkers using the Centers for Disease Control and Prevention (CDC) categories to define "[underweight](#)," "normal weight," "overweight" and "obesity."

"The NHIS is like a 'selfie' for the U.S. because it is a snapshot of health behaviors of [people](#) from every type of background," said Muntasir Masum, postdoctoral scholar at the Edna Bennett Pierce Prevention Research Center at Penn State. "We expected to see a link between obesity and mortality related to alcoholism, and we were surprised to see that the link was especially pronounced for people with underweight who drink excessively."

The CDC defines underweight as having a [body mass index](#) (BMI) of less than 18.5 using the calculation of person's weight in kilograms divided by the square of their height in meters. According to the CDC, BMI is a screening tool, but it does not diagnose "body fatness or the health of an individual."

Further research is needed into how having underweight could contribute to mortality in people who drink excessively. Masum suggested that multiple factors could be at play, such as how people handle stress and whether they have co-occurring health issues or nutritional deficiencies.

"I hope these findings encourage people to eliminate risks that may lead to a life-or-death situation," said Masum.

Excessive alcohol use is the third most common cause of preventable death in the U.S. and is estimated to cause 1 in 10 deaths among working-age adults in the U.S., according to the *Journal of the American Medical Association*.

This work was supported by Penn State's Prevention and Methodology Training program, which is funded by the National Institute on Drug Abuse.

Collaborators on the study included Jeffrey Howard, associate professor of public [health](#) at the University of Texas at San Antonio; and Timothy Grigsby, assistant professor in the Department of Social and Behavioral Health at the University of Nevada, Las Vegas.

**More information:** Muntasir Masum et al, Combined alcohol use and weight status effects on mortality risk among adults: Analysis of the National Health Interview Survey linked mortality files, 2001–2015, *Drug and Alcohol Dependence* (2021). [DOI: 10.1016/j.drugalcdep.2021.109206](#)

Avani K. Patel et al, Unhealthy Alcohol Use, *JAMA* (2021). [DOI: 10.1001/jama.2020.2015](#)

Provided by Pennsylvania State University

Citation: How can body weight affect the mortality risk of excessive drinkers? (2022, January 24) retrieved 19 April 2024 from <https://medicalxpress.com/news/2022-01-body-weight-affect-mortality-excessive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.