

Live cells discovered in human breast milk could aid breast cancer research

January 28 2022



Credit: Pixabay/CC0 Public Domain

Researchers have explored the cellular changes that occur in human mammary tissue in lactating and non-lactating women, offering insight into the relationship between pregnancy, lactation, and breast cancer.

The study was led by researchers from the Wellcome-MRC Cambridge Stem Cell Institute (CSCI) and the Department of Pharmacology at the University of Cambridge.

Breast tissue is dynamic, changing over time during puberty, pregnancy, breastfeeding, and aging. The paper, published today in the journal *Nature Communications*, focuses on the changes that take place during lactation by investigating [cells](#) found in [human milk](#).

This research, led by Dr. Alecia-Jane Twigger of CSCI, found that the cells in [milk](#), once thought to be dead or dying, are in fact very much alive. These [living cells](#) provide researchers with the chance to study not only the changes that occur in mammary tissues during lactation, but also insight into a potential early indicator of future breast cancer development.

"I believe that by studying human milk cells, we will be able to answer some of the most fundamental questions around mammary gland function such as: how is milk produced? Why do some women struggle to make milk? and what strategies can be employed to improve breastfeeding outcomes for women?" said Dr. Alecia-Jane Twigger at the Wellcome-MRC Cambridge Stem Cell Institute who led the study.

The researchers collected voluntary breast milk samples from lactating women, as well as samples of non-lactating breast tissue donated from women who elected to have esthetic breast reduction surgery. Using single-cell RNA sequencing analysis, the team conducted a novel comparison of the composition of the mammary cells taken using these two methods, identifying the distinctions between lactating and non-lactating human mammary glands.

While accessing breast tissue for study relies on donors already undergoing surgery, breast milk samples are much simpler to acquire.

Breast milk donors are engaged via midwives or women's networks (an undertaking made more challenging by the pandemic) and agree to share their samples over time. Typical daily production for lactating women is between 750-800ml, and the sample size for Twigger's research is on average a mere 50ml, an amount which can contain hundreds of thousands of cells for study.

By collecting these samples donated by breastfeeding women—samples now known to contain living and viable cells—researchers have the opportunity to capture dynamic cells in a non-invasive way. This greater ease of access to [breast](#) cells can open the door to more studies on women's health in the future.

"The first time Alecia told me that she found [live cells](#) in milk I was surprised and excited about the possibilities. We hope this finding will enable future studies into the early steps of [breast cancer](#)," said Dr. Walid Khaled, at the Wellcome-MRC Cambridge Stem Cell Institute and University of Cambridge's Department of Pharmacology, who was also involved in the study.

More information: Transcriptional changes in the mammary gland during lactation revealed by single cell sequencing of cells from human milk, *Nature Communications* (2022). [DOI: 10.1038/s41467-021-27895-0](#)

Provided by University of Cambridge

Citation: Live cells discovered in human breast milk could aid breast cancer research (2022, January 28) retrieved 25 April 2024 from <https://medicalxpress.com/news/2022-01-cells-human-breast-aid-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.