

To reduce COVID-19 risk, individual distancing makes a difference

January 24 2022, by Beth Newcomb



Credit: Unsplash/CC0 Public Domain

As the COVID-19 pandemic stretches into 2022 and vaccination campaigns continue across the world, individual health behaviors are still a key factor in the risk of COVID-19 diagnosis, according to a new USC

study.

A new paper by USC Leonard Davis School of Gerontology researchers illustrates how an individual's physical distancing behaviors, such as avoiding large gatherings, can significantly decrease the [risk](#) of COVID-19 diagnosis, even in the presence of other structural risk factors, such as high local infection rates.

"Personal mitigation behaviors related to [social distancing](#) appear to influence risk of COVID-19 even in the presence of social factors related to infection risk," said Theresa Andrasfay, the study's first author and postdoctoral scholar at the USC Leonard Davis School. "Our findings emphasize the importance of individual behaviors for preventing COVID-19, which may be relevant in contexts with low vaccination."

The study examined responses from the COVID-19 survey of the Understanding America Study from June 2020 through April 2021. The UAS includes a panel of approximately 9,500 respondents representing the entire United States and is administered by the USC Center for Economic and Social Research.

Andrasfay and colleagues focused on respondents' answers to questions about whether they had been diagnosed with COVID-19 and if they have taken part in 11 risky behaviors, from going out to bars or restaurants to hosting non-household members at one's residence. The team was able to adjust the data based on sex, age, race/ethnicity, education, employment status, living arrangements, as well as the statewide 7-day COVID-19 case rate at the time of responses.

The researchers found that individuals who reported attending large gatherings were more than 40% more likely to have been diagnosed with COVID-19 than those who did not report unsafe behaviors, even after

adjusting for demographic factors and local case rates. In respondents who took part in more than one of the risky behaviors included in the survey, each additional risk taken increased the likelihood of COVID-19 diagnosis by another 8 to 9%.

Even among those who avoided risks involving large gatherings, taking part in smaller gatherings still increased the risk of diagnosis by 30%. This is consistent with earlier research suggesting that small birthday gatherings were associated with household COVID-19 infections, Andrasfay noted.

The survey didn't include detailed measures about mask wearing, such as type of mask worn and how diligently one wore it, which limited the researcher's ability to detect the protective effect of mask wearing in the study. "However, we know from many other studies that high quality masks are one of the best ways to prevent COVID-19," Andrasfay said.

Even with research into treatments and a push for widespread vaccination, the study highlights the continuing importance of individuals reducing their risk through [behavior](#), especially physical distancing and avoiding large gatherings, she said.

"For much of the COVID-19 pandemic, nonpharmaceutical interventions, including behavioral modifications, were the only defense against COVID-19," Andrasfay said. "Although vaccines are now widely available in the United States, the surges in infections fueled by the Delta and Omicron variants underscores that behavioral modifications continue to be important for reducing the spread of COVID-19."

"Adherence to Social-Distancing and Personal Hygiene Behavior Guidelines and Risk of COVID-19 Diagnosis: Evidence From the Understanding America Study" first appeared online in the *American Journal of Public Health* on December 22, 2021.

More information: Theresa Andrasfay et al, Adherence to Social-Distancing and Personal Hygiene Behavior Guidelines and Risk of COVID-19 Diagnosis: Evidence From the Understanding America Study, *American Journal of Public Health* (2021). [DOI: 10.2105/AJPH.2021.306565](https://doi.org/10.2105/AJPH.2021.306565)

Provided by University of Southern California

Citation: To reduce COVID-19 risk, individual distancing makes a difference (2022, January 24) retrieved 8 May 2024 from <https://medicalxpress.com/news/2022-01-covid-individual-distancing-difference.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.