

Hip fracture surgery generally lower during COVID-19 pandemic

January 4 2022



(HealthDay)—Among older individuals, hip fracture surgery volumes



were generally lower during the COVID-19 pandemic than two years earlier until the COVID-19 vaccine became available, according to a research letter published online Dec. 30 in the *Journal of the American Medical Association*.

Kanu Okike, M.D., M.P.H., from Kaiser Permanente in Honolulu, and colleagues examined hip fracture surgery volumes among individuals ages 65 years or older during the pandemic. The incidence rate was calculated for each of eight phases during the pandemic (presurge [Jan. 1 to March 15, 2020], spring 2020 surge [March 16 to May 10], early summer 2020 surge [May 11 to June 28], summer 2020 surge [June 29 to Aug. 23], fall 2020 surge [Aug. 24 to Nov. 22], winter surge [Nov. 23, 2020, to Feb. 21, 2021], early vaccination period [Feb. 22 to July 18, 2021], and delta wave [July 19 to Sept. 26, 2021]) and were compared to the historical period two years prior.

The researchers found that during the presurge period, the <u>volume</u> of hip fracture surgery was similar to that of the historical period, but was lower in the spring 2020 surge (incidence rate ratio, 0.81; 95 percent confidence interval, 0.73 to 0.90; P period (incidence rate ratio, 0.96; 95 percent confidence interval, 0.90 to 1.02; P = 0.18) and the delta wave.

"Further research should determine how volume patterns evolve with increasing vaccinations and COVID-19 variants and investigate the reasons for the lower incidence, which may provide insight into ways to decrease hip <u>fractures</u> among older individuals," the authors write.

More information: Abstract/Full Text

Copyright © 2021 HealthDay. All rights reserved.

Citation: Hip fracture surgery generally lower during COVID-19 pandemic (2022, January 4)



retrieved 4 May 2024 from https://medicalxpress.com/news/2022-01-hip-fracture-surgery-covid-pandemic.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.