

The impacts of local iodine variations on human health

January 17 2022, by Thea Cecilie Engelsen



Iodine and health in Somaliland



Is iodine intake sufficient in Somalilanders?

The findings in the iodine surveys during the last ten years are conflicting. But they can all be correct: In some regions, people may have iodine deficiency, in others, iodine excess! Either condition increaseas the risk of low thyroid hormon production

variation is large. Average lodine Concentration (µg/L)



Journal of Nutrition https://doi.org/10.1093/jn/nxob377

Research has shown that ground water in Somaliland contains high levels of iodine. The geographical

Half of the iodine intake originates from drinking water, the rest from food.



Dry hair and nails Poor appetite Constipation Muscle weakness e.k.been@medisin.uio.no, a.a.madar@medisin.uio.no

What should be done?

Measure iodine concentrations in all main water sources throughout a complete year.

Investigate iodine status in villages with high and low water iodine, including breastfeeding women and small children.

Use the findings for targeted public health interventions.

Credit: University of Oslo



Ground water is the main source of iodine to the Somali population, but the geographical variation is considerable. While earlier studies from the region have concluded the iodine intake to be excessive or that there is general iodine deficiency in the Somali population, researcher Espen Heen, and his team, show that both may be correct.

"In some areas the ground water hold high concentrations of iodine while in other areas it is more moderate or low," says Heen, who worries that general measures to improve the intake of iodine, if implemented for the population as a whole, will affect the Somali people negatively.

Too little iodine during pregnancy affects the development of the brain and can cause severe sensory and motor impairments as well as reduced IQ. Further, in children and adults, it can affect the thyroid and cause hypothyroidism—a condition in which the <u>thyroid gland</u> does not produce enough of certain crucial hormones. Too much iodine can also damage the thyroid gland and lead to hyperthyroidism with subsequent heart failure.

Adding iodine to salt

"A good balance of iodine is crucial to a person's health. But with the correct measures it is a relatively easy public health issue to solve," says Heen.

The most common way to improve a population's iodine intake is by adding iodine to salt. This is also the <u>recommended strategy from WHO</u>.

While this strategy may be efficient in a population with relatively similar iodine exposal, Heen and his team believe that their study, newly published in *Journal of Nutrition*, shows that this is not the case in Somaliland. They also believe their study describes a situation that is relevant for neighboring countries on the Horn of Africa and other



countries with similar variations in iodine intake.

"This study shows how we need to rethink the methodology of assessing and introducing iodine to large population groups in such areas. Locally adapted measures are vital to balance the population's intake of iodine. Even across small geographical areas we detected notable differences in ground water iodine levels."

Twofold study

The study that Heen and his team has conducted is twofold. They have compared standard and new assessment methodology in samples from 118 non-pregnant, healthy women aged 15–69 years residing in Hargeisa—the capital of Somaliland in the northern part of Somalia. 27 of the women were breastfeeding. The researchers have also gone back to two previous studies in Somalia and re-assessed these results based on their recent findings.

It is the first time the correlation between ground water iodine and iodine intake in the region is established firmly.

"We measured both the urinary iodine concentration and the household water iodine concentration across 24 hours. There were strong correlations between these, with iodine from household <u>water</u> contributing more than one-half of the total iodine intake. Physiological hydration and breastfeeding status we found to be the main predictors of urinary <u>iodine</u> concentration."

Heen has conducted his research together with colleagues at the Institute of Health and Society at the University of Oslo in collaboration with Somali health workers and researchers at The University of Hargeisa. The project was part of a larger international NORAD-funded capacitybuilding project in Somalia.



The next step for Heen and his team is now to reach a dialog with local health authorities in Somaliland and with UN organization about how to follow up and assess the findings further.

More information: Espen Heen et al, Household Water is the Main Source of Iodine Consumption Among Women in Hargeisa, Somaliland: A Cross-Sectional Study, *The Journal of Nutrition* (2021). DOI: <u>10.1093/jn/nxab377</u>

Provided by University of Oslo

Citation: The impacts of local iodine variations on human health (2022, January 17) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2022-01-impacts-local-iodine-variations-human.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.