

Study examines years of life lost for adolescents, young adults to drug overdose

January 31 2022



Credit: CC0 Public Domain

In a first-ever study, researchers at The Ohio State University Wexner Medical Center and College of Medicine examined the unintentional drug overdose mortality in years of life lost among adolescents and

young people in the United States.

Study findings are published online in the journal *JAMA Pediatrics*.

Excess mortality was calculated in Years of Life Lost (YLL), which is the difference between the age at which a person dies and their expected remaining lifespan. In 2019, U.S. life expectancy at birth was 78.8 years, according to the Centers for Disease Control and Prevention.

"Adolescents and [young adults](#) dying of [overdose](#) are deprived of many years of work, community, and [family life](#). Our study shows overdose mortality among adolescents and young people is unacceptably high. Public health interventions to protect this vulnerable group are urgently needed," said Dr. O. Trent Hall, first author of the study and an addiction medicine physician in Ohio State's Department of Psychiatry and Behavioral Health.

This study looked at 3,296 adolescents ages 10-19 and 21,689 young people ages 10-24 who lost their lives from unintentional [drug overdose](#) from 2015-2019. During this time, adolescents lost nearly 200,000 years of life due to unintentional [drug](#) overdose, while young people amassed more than 1.25 million years of lost life from overdose.

Even though adolescents and young people are dying in increasing numbers from drug overdose, adults have been the primary focus of most research reports, Hall said.

"We hope our report will raise awareness of the increasingly dire consequences of unintentional drug overdose among this vulnerable population," said Dr. Julie Teater, senior author on the study and a psychiatrist and addiction medicine physician in Ohio State's Department of Psychiatry and Behavioral Health. "Our study provides important context to the overdose crisis by better representing what it

means to society when we lose adolescents and young people to unintentional drug overdose."

This study builds on Hall's prior research that had calculated years of life lost due to unintentional drug overdose in Ohio among the general population and pregnant individuals or those within one year of pregnancy.

"A clear implication of our research is that more resources are needed to prevent unintentional drug overdose among adolescents and [young people](#). Existing [public health interventions](#) aimed at adults may be insufficient," Hall said.

More research is needed to determine how best to engage younger individuals with education, prevention, harm reduction, substance treatment, as well as community and family support. This study points to the need to improve interventions involving family, school and community in drug overdose prevention and substance use treatment, Teater said.

Provided by Ohio State University Medical Center

Citation: Study examines years of life lost for adolescents, young adults to drug overdose (2022, January 31) retrieved 20 April 2024 from <https://medicalxpress.com/news/2022-01-years-life-lost-adolescents-young.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--