


If you're over 65, you're more likely to die with kidney disease than from it

January 18 2022



Credit: Robina Weermeijer/Unsplash

People over 65 years of age with stage 4 chronic kidney disease (CKD) were more likely to die than to experience kidney failure, suggesting that more people die with CKD rather than directly because of CKD, according to research published today by the *Medical Journal of Australia*.

Professor Matthew Jose, Professor of Medicine at the University of Tasmania and Royal Hobart Hospital, and colleagues analyzed linked data from the Tasmanian Chronic Kidney Disease study, for all Tasmanian adults diagnosed with incident stage 4 [chronic kidney disease](#) between 1 January 2004 and 31 December 2017. Data for a total of 6825 adults (mean age, 79.3 years; SD, 11.1 years), including 3,816 women (55.9 percent), was analyzed.

"The risk of death increased with age—under 65 years: 0.18 (95 percent CI, 0.15–0.22); 65–74 years: 0.39 (95 percent CI, 0.36–0.42); 75–84 years, 0.56 (95 percent CI, 0.54–0.58); 85 years or older: 0.78 (95 percent CI, 0.77–0.80)—while that of [kidney](#) failure declined—under 65 years: 0.39 (95 percent CI, 0.35–0.43); 65–74 years: 0.12 (95 percent CI, 0.10–0.14); 75–84 years: 0.05 (95 percent CI, 0.04–0.06); 85 years or older: 0.01 (95 percent CI, 0.01–0.02)," the authors reported.

"Three important conclusions can be drawn from our study: for [people](#) with stage 4 CKD, the 5-year risk of kidney failure declines with age; the [risk](#) of death is greater than that of kidney failure for those over 65 years of age; and the risks of death and kidney failure for people over 65 are each greater for men than women.

"People over 65 years of age with stage 4 CKD were more likely to die than to experience kidney failure, suggesting that more people die with CKD rather than directly because of CKD," Jose and colleagues wrote.

"When kidney replacement therapy pathways are well defined and supported, disease management typically focuses on dialysis or transplantation; the pathway to supportive (non-dialysis) care is less well defined, and differences between renal units are greater.

"[...] mortality risks should be considered when discussing future kidney failure treatment options for patients over 65 years of age."

Currently [clinical guidelines](#) recommend that patients be referred to kidney specialists, in order to prepare for [kidney failure](#) or kidney replacement therapy.

Jose and colleagues wrote that their findings showed that "clinical guidelines should recognize competing risks of death in [older people](#) with severe kidney disease, and consider holistic, supportive care, not just kidney replacement therapy."

More information: Matthew D Jose et al, Competing risks of death and kidney failure in a cohort of Australian adults with severe chronic kidney disease, *Medical Journal of Australia* (2021). [DOI: 10.5694/mja2.51361](#)

Provided by Medical Journal of Australia

Citation: If you're over 65, you're more likely to die with kidney disease than from it (2022, January 18) retrieved 5 May 2024 from <https://medicalxpress.com/news/2022-01-youre-die-kidney-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--