

# Bilateral oophorectomy could increase a woman's risk for dementia

February 2 2022

---



Credit: Pixabay/CC0 Public Domain

Despite the advantage of an oophorectomy reducing the threat of ovarian cancer, it is not without risk. Removing the ovaries causes premature menopause and hormone disruption that can lead to heart disease, osteoporosis, depression, and other problems. A new study adds to the literature and shows an increased risk of dementia with oophorectomy.

Study results are published online today in *Menopause*.

Women are disproportionately affected by dementia, with a 55% higher risk from age 65 years onward, compared with men. This sex difference is not fully explained by the fact that [women](#), on average, live longer than men.

Estrogen has been consistently identified as a potential mechanism underlying higher rates of dementia in women. That's why common interventions that permanently alter sex steroid exposure, such as oophorectomy, are suspected of representing a female-specific risk factor of dementia. Bilateral oophorectomy (the removal of both ovaries) is sometimes performed in conjunction with hysterectomy to help prevent ovarian cancer in women with a high inherited risk.

Several earlier studies investigated the association between [bilateral oophorectomy](#) and dementia but showed inconsistent findings. In this new study involving nearly 25,000 women, the goal was not to simply investigate the prospective association between oophorectomy and the rate of incident dementia but also to determine whether the risk was affected by age at the time of surgery, hysterectomy, or use of hormone therapy.

On the basis of their results, the researchers concluded that bilateral, but not unilateral, oophorectomy was associated with an increased risk of dementia. However, they were not able to show whether this association was conditional with hysterectomy or hormone therapy use.

Study results are published in the article "Oophorectomy and rate of dementia: a prospective cohort study."

"These results are consistent with findings of prior studies showing a link between oophorectomy and [dementia](#). The statistical power of the study

was limited, which may explain the lack of effect of age or [hormone](#) therapy use on this association. Given the robust body of evidence suggesting potential long-term adverse effects associated with oophorectomy before the average age of menopause, risk-reducing oophorectomy should be limited to those women with an inherited high risk for cancer," says Dr. Stephanie Faubion, NAMS medical director.

**More information:** Cecilie S. Uldbjerg et al, Oophorectomy and rate of dementia, *Menopause* (2022). [DOI: 10.1097/GME.0000000000001943](#)

Provided by The North American Menopause Society

Citation: Bilateral oophorectomy could increase a woman's risk for dementia (2022, February 2) retrieved 26 June 2024 from <https://medicalxpress.com/news/2022-02-bilateral-oophorectomy-woman-dementia.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--