

Researchers find new clues in the brain linking pain and food

February 11 2022, by Kelsie Smith Hayduk



Credit: Pixabay/CC0 Public Domain

It has long been known that there is an association between food and pain, as people with chronic pain often struggle with their weight. Researchers at the Del Monte Institute for Neuroscience may have found

an explanation in a new study that suggests that circuitry in the brain responsible for motivation and pleasure is impacted when someone experiences pain. "These findings may reveal new physiological mechanisms linking chronic pain to a change in someone's eating behavior," said Paul Geha, M.D., lead author on the study published in *PLOS ONE*. "And this change can lead to the development of obesity."

Finding pleasure in food comes from how our [brain](#) responds to what we are eating. In this study researchers were looking at the brain's response to sugar and fat. Using a gelatin dessert and pudding, researchers altered the sugar, fat, and texture of the foods. They found that none of the patients experienced eating [behavior](#) changes with sugar, but they did with fat. Those with acute lower back pain who later recovered were most likely to lose pleasure in eating the pudding and show disrupted satiety signals—the communication from the digestive system to the brain—while those with acute lower back pain whose pain persisted at one year did not initially have the same change in their eating behavior. But chronic lower back pain patients did report that eventually foods high in fat and carbohydrates, like [ice cream](#) and cookies, became problematic for them over time and brain scans showed disrupted satiety signals.

"It is important to note, this change in food liking did not change their [caloric intake](#)," said Geha, who first authored a previous study published in *PAIN* that recent research is building on. "These findings suggest obesity in patients with chronic pain may not be caused by lack of movement but maybe they change how they eat."

Brain scans of the study participants revealed that the nucleus accumbens—a small area of the brain mostly known for its role in decision-making—may offer clues to who is at risk to experience a long-term change in eating behavior. Researchers found the structure of this area of the brain was normal in of patients who initially experienced

changes in their eating behavior but whose pain did not become chronic. However, patients whose eating behavior was normal, but whose pain became chronic had smaller nucleus accumbens. Interestingly, the nucleus accumbens predicted pleasure ratings only in chronic back pain patients and in patients who became chronic after an acute bout of back pain suggesting that this region becomes critical in motivated behavior of chronic [pain](#) patients. Previous research by Geha, found a smaller [nucleus accumbens](#) can indicate if someone is at a greater risk of developing [chronic pain](#).

Additional authors include Yezhe Lin, Ph.D., and Gelsina Stanley of the University of Rochester, Ivan de Araujo, Ph.D., of Icahn School of Medicine at Mount Sinai, and Dana Small, Ph.D., of Yale University.

More information: Yezhe Lin et al, Chronic pain precedes disrupted eating behavior in low-back pain patients, *PLOS ONE* (2022). [DOI: 10.1371/journal.pone.0263527](#)

Provided by University of Rochester Medical Center

Citation: Researchers find new clues in the brain linking pain and food (2022, February 11) retrieved 11 July 2024 from <https://medicalxpress.com/news/2022-02-clues-brain-linking-pain-food.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
