


COVID-19 boosters and some medications appear to neutralize omicron

February 3 2022



Credit: Dr Jason A. Roberts, head, Electron Microscopy and Structural Virology, Doherty Institute

Mutations to the spike protein of the omicron variant are shown to alter how the virus infects cells and reduce its sensitivity to therapeutic and vaccine-elicited antibodies, according to a paper published in *Nature*. Booster doses of mRNA vaccines improve neutralization, and the antiviral drugs remdessevir and molnupiravir are shown to retain efficacy

against omicron.

[omicron](#) carries approximately 36 mutations in the spike protein, the part of the virus that facilitates entry into host [cells](#). Ravindra Gupta and colleagues identify changes that have the potential to affect binding to the host receptor, ACE2. They also show altered infectivity in several [cell types](#), including reduced replication in airway organoids and gut cells, compared with the previous delta variant. They show that omicron is less dependent on the cell membrane protein transmembrane serine protease 2 (TMPRSS2), which is required by previous variants for efficient infection of cells. Their data indicate that mutations to the omicron spike reduce efficient infection of lower airway cells expressing TMPRSS2 (such as cells in the alveoli), but not TMPRSS2-negative cells such as those found in the upper airway.

To understand how omicron may respond to existing COVID-19 treatments, the authors incubated both live and pseudoviral particles with clinically approved antibodies to analyze the interactions between the antibodies and the omicron [spike](#). They find that casivirimab and imdevimab, which have been shown to be particularly effective at neutralizing delta when used together, lose all neutralizing activity against omicron when used either individually or in combination. The antiviral agents remdesivir and molnupiravir are shown to have similar activity against delta and omicron.

To determine the effectiveness of existing vaccines against omicron, Gupta and co-workers produced [virus particles](#) expressing the [spike protein](#) from omicron or delta, and exposed them to serum from individuals vaccinated with either the Pfizer–BioNTech or Oxford–AstraZeneca vaccine. They observed at least a 10-fold reduction in neutralization of omicron compared to delta after two doses of either vaccine, with further waning in response associated with increasing time since the second dose. Neutralization of omicron was not detectable for

the majority of individuals who had received two doses of Oxford–AstraZeneca. Further sera from the same groups, but boosted with Pfizer–BioNTech as a third dose, was shown to increase neutralization. Similar results were observed for two doses of Pfizer–BioNTech in experiments using live virus.

More information: Bo Meng et al, Altered TMPRSS2 usage by SARS-CoV-2 Omicron impacts tropism and fusogenicity, *Nature* (2022). [DOI: 10.1038/s41586-022-04474-x](https://doi.org/10.1038/s41586-022-04474-x)

Provided by Nature Publishing Group

Citation: COVID-19 boosters and some medications appear to neutralize omicron (2022, February 3) retrieved 1 May 2024 from <https://medicalxpress.com/news/2022-02-covid-boosters-medications-neutralize-omicron.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
