

Feeling dizzy when you stand up? Simple muscle techniques can effectively manage symptoms

February 9 2022



Investigators determined both lower body muscle preactivation (PREACT) and lower body muscle tensing (TENSE) effectively reduced the drop in mean arterial pressure upon standing and improved symptoms of initial orthostatic hypotension. Left: Schematic diagram of the three interventions: no intervention (Control), PREACT, and TENSE. Right: Drop in mean arterial pressure following Control (blue), PREACT (pink), and TENSE (purple). Credit: Heart Rhythm



Feeling lightheaded upon standing up due to initial orthostatic hypotension (IOH), or a transient decrease in blood pressure and increase in heart rate, is a common but poorly understood condition. A new study offers two simple cost- and drug-free techniques to effectively manage symptoms of IOH and improve quality of life by activating lower body muscle before or after standing. The research appears in *Heart Rhythm*, the official journal of the Heart Rhythm Society, the Cardiac Electrophysiology Society, and the Pediatric & Congenital Electrophysiology Society.

Syncope, lightheadedness, dizziness, or loss of consciousness from IOH, affects up to 40% of the general population (all ages), while presyncope is probably even more common. Despite this, the condition is relatively understudied and there is minimal information available about the underlying mechanisms or symptom management and treatment. Currently, there are very few options available to patients with IOH and no pharmacological treatments. The most common recommendations have been to stand up slowly or sit up first before standing.

"Almost everyone has probably experienced some lightheadedness at some time after standing up," explained lead investigator Satish R. Raj, MD, MSCI, FHRS, Professor of Cardiac Sciences, Libin Cardiovascular Institute, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada. "For some people this is a frequent occurrence and may happen several times a day, which can be very frightening and negatively impact their quality of life. We wanted to explore this further and provide novel and effective symptom management techniques with the goal of improving the IOH patient's quality of life."

This study investigated physical maneuvers before or after standing and their efficacy in reducing the drop in <u>blood pressure</u> as well as the symptoms typically seen in IOH patients upon standing. Study participants included 24 young women (mean age 32 ± 8 years) with a



high burden history of fainting immediately after standing and more than four episodes of presyncope or syncope per month. The study participants were required to have a significant drop in <u>systolic blood</u> <u>pressure</u> of at least 40 mmHg upon standing to fulfill the diagnostic criteria of IOH on the study day. Two participants had inadequate <u>heart</u> <u>rate</u> recordings and were excluded from the analysis.

The 22 study participants completed three sit-to-stand maneuvers including a stand with no intervention (control), and two interventions. Researchers found that both lower body muscle preactivation (thighs) through repeated knee raises prior to standing (PREACT) and lower body muscle tensing (thighs and buttocks) through leg crossing and tensing immediately after standing (TENSE) effectively improve the blood pressure drop. This led to a reduction in symptoms upon standing. They found that the PREACT maneuver accomplished this by increasing cardiac output, while the TENSE maneuver did so by increasing stroke volume.

"Our study provides a novel and cost-free symptom management technique that patients with IOH can use to manage their symptoms," noted first author Nasia A. Sheikh, MSc, Libin Cardiovascular Institute, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada. "Since it is a physical maneuver, it simply requires the lower body limbs, which patients can utilize at any time and from anywhere to combat their symptoms."

"Our study demonstrates the physiology of IOH and assesses the utility of physical maneuvers that can help the IOH patient manage their symptoms. A diagnosis of IOH is identified by patients as the critical first step to empowering them to understand and master their symptoms and thus minimize the disruptions to daily living caused by this common, but not commonly understood, condition," added co-investigator Mary Runté, Ph.D., University of Lethbridge, Lethbridge, AB, Canada.



In an accompanying editorial, Bert Vandenberk, MD, Ph.D., Department of Cardiac Sciences, Libin Cardiovascular Institute, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada; and Department of Cardiovascular Sciences, KU Leuven, Leuven, Belgium; and Carlos A. Morillo, MD, FHRS, Department of Cardiac Sciences, Libin Cardiovascular Institute, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada, commented that "PREACT and TENSE provide an elegant and simple effective <u>symptom</u> relief option for patients with IOH." However, they noted, "The study needs to be validated in men, as well as being explored in the elderly where multiple confounders should be addressed. Understanding the role of cardiopulmonary mechanoreceptors should also provide further mechanistic insights into the effects of these simple yet very effective physiologic maneuvers."

More information: Nasia A. Sheikh et al, Lower body muscle preactivation and tensing mitigate symptoms of initial orthostatic hypotension in young females, *Heart Rhythm* (2022). <u>DOI:</u> 10.1016/j.hrthm.2021.12.030

Bert Vandenberk et al, Symptom reduction in initial orthostatic hypotension: Time to get physical!, *Heart Rhythm* (2022). DOI: 10.1016/j.hrthm.2021.12.031

Provided by Elsevier

Citation: Feeling dizzy when you stand up? Simple muscle techniques can effectively manage symptoms (2022, February 9) retrieved 7 May 2024 from https://medicalxpress.com/news/2022-02-dizzy-simple-muscle-techniques-effectively.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.