


# Early life treatment prevents autism symptoms from developing in mice

February 21 2022



Treatment with rapamycin for 4 weeks prevents the onset of social deficits in three-chamber social approach (left) and social novelty (right) testing. Credit: Gibson et al., *JNeurosci* 2021

Timing is key when treating developmental disorders. Blocking an overactive signaling pathway during the first five weeks of life prevents autism symptoms from ever developing in mice, according to new research published in *JNeurosci*.

The brain develops capabilities, like language, during specific spans of time called critical periods. The symptoms of disorders like autism arise during a set critical period; administering a targeted intervention only during the critical period could prevent the disorder from ever taking shape without the burden of a life-long treatment.

Gibson et al. compared intervention timing in a genetic mouse model of Tuberous Sclerosis Complex (TSC), a [neurodevelopmental disorder](#) with high rates of autism spectrum disorder. The research team administered the drug rapamycin to inhibit mTOR signaling, a pathway that is overactive in people and animals with TSC. After four weeks of treatment starting at one week of life, [mutant mice](#) displayed normal social behaviors and normal activity in cerebellar Purkinje neurons, even after four weeks without treatment.

These results raise the possibility that treatment covering critical periods of development could prevent [autism symptoms](#) from ever manifesting. As TSC is often diagnosed *in utero* in humans, defining these critical periods may provide an opportunity to prevent development of autism behaviors.

**More information:** A Critical Period for Development of Cerebellar-Mediated Autism-Relevant Social Behavior, *JNeurosci* (2022). [DOI: 10.1523/JNEUROSCI.1230-21.2021](#)

Provided by Society for Neuroscience

Citation: Early life treatment prevents autism symptoms from developing in mice (2022, February 21) retrieved 11 July 2024 from <https://medicalxpress.com/news/2022-02-early-life-treatment-autism-symptoms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.