

More than half of postmenopausal women experience female pattern hair loss

February 16 2022



Credit: Laura Tiitto/public domain

Both men and women are more likely to lose their hair with age. Women also have the added risk associated with declines in estrogen levels during the menopause transition. A new study sought to identify the prevalence of female pattern hair loss (FPHL), hair characteristics, and associated factors in healthy postmenopausal women. Study results are

published online today in *Menopause*.

Female pattern [hair](#) loss is the most common hair loss disorder in women. It is characterized by gradual thinning at the part line, followed by increasing diffuse hair loss radiating from the top of the head. Female pattern hair loss can develop any time between the teenage years and the postmenopause period. However, it is believed that the loss of estrogen during the menopause transition may play a role in accelerating FPHL because estrogen receptors are present in hair follicles. Menopause-related hormone changes have been shown to influence scalp hair, reduce hair diameter, and limit hair growth.

Hair loss can have a significant effect on a woman's self-esteem and overall quality of life because it affects her appearance and confidence. Because women spend, on average, one third of their lives postmenopause, research into causes and treatments of hair loss is critical.

In a new cross-sectional study involving 178 women seen at a menopause clinic, researchers aimed to evaluate the prevalence of FPHL in healthy postmenopausal women and investigate postmenopausal hair characteristics as well as the factors associated with FPHL. Of the women studied, 52.2% were found to have FPHL. The prevalence of FPHL increased with age. Low self-esteem was detected in 60% of participants and increased with the severity of FPHL.

The researchers additionally noted that a high body mass index (obesity) was associated with an increased prevalence and worsening of FPHL in postmenopausal women. Further studies are necessary to determine whether sex steroid hormones, especially estrogen and testosterone, and a history of polycystic ovary syndrome are related to hair loss in postmenopausal women.

Study results are published in the article "Prevalence of female pattern hair loss in postmenopausal women: a cross-sectional study."

"Female pattern hair loss was prevalent and associated with lower [self-esteem](#) in healthy [postmenopausal women](#) in this small cross-sectional study. A better understanding of the mechanisms responsible for this common type of hair loss in [women](#) may lead to more effective preventive strategies and treatment options," says Dr. Stephanie Faubion, NAMS medical director.

More information: Prevalence of female pattern hair loss in postmenopausal women: a cross-sectional study, *Menopause* (2022). [DOI: 10.1097/GME.0000000000001927](https://doi.org/10.1097/GME.0000000000001927)

Provided by The North American Menopause Society

Citation: More than half of postmenopausal women experience female pattern hair loss (2022, February 16) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-02-postmenopausal-women-female-pattern-hair.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--