

Study finds potential predictor for perimenopause

February 11 2022



Credit: Pixabay/CC0 Public Domain

A new study has found women 'age' significantly faster in the perimenopausal period, based on analysis of the IgG glycome. This finding indicates the potential of using the IgG glycome composition as

an early predictor for perimenopause.

The study, published today in *iScience*, and led by team of researchers from the Genos Glycoscience Research Laboratory in Zagreb, Croatia, Newson Health Menopause & Wellbeing in Stratford upon Avon and King's College London followed nearly 2000 women for 15 years and analyzed their IgG glycans several times during that period. IgG glycans, otherwise known as immunoglobulin G glycome, is an abundant antibody and has been shown to be a biomarker of an individual's health.

Researchers analyzed IgG glycome composition in 5,080 samples from 1,940 females during their transition from before menopause (premenopause) to postmenopause. Many of these samples were taken from TwinsUK which is the largest twin registry in the UK and led by researchers from King's College London.

Analysis of the IgG glycome in multiple samples from the same individuals was able to show the association between perimenopause and the changes in the IgG glycome composition. This period showed the IgG glycome changing from an inflammation-suppressive to proinflammatory. While this is common as we age, researchers saw this change happen rapidly as women transitioned from a regular cycle to menopause.

This change in IgG glycans is associated with many health risks that accompany menopause. In some diseases like [rheumatoid arthritis](#) and cardiovascular diseases, this change can occur years before disease onset. This suggest glycans and [chronic inflammation](#) is linked and the transformation of IgG glycans plays a part in developing disease.

Perimenopause can last up to 15 years and is challenging to diagnose due to highly irregular hormonal cycles. As a result of poor awareness and inappropriate use of hormonal tests, women are often misdiagnosed with

conditions such as fibromyalgia, migraines, depression, or [chronic fatigue syndrome](#) and are frequently prescribed antidepressants despite the lack of evidence to support their use to improve the low mood associated with perimenopause or menopause. This study shows it may be possible to use IgG glycans to predict perimenopause.

Dr. Cristina Menni, from TwinsUK and King's College London, said that "Perimenopause is poorly diagnosed due to highly irregular hormonal cycles and symptoms that can last for as long as 15 years. Currently, there is no accurate diagnostic test for perimenopause. Adding an easily quantifiable novel early biomarker of [perimenopause](#) could be a valuable improvement to current clinical praxis."

More information: Helena Deriš et al, Immunoglobulin G glycome composition in transition from premenopause to postmenopause, *iScience* (2022). [DOI: 10.1016/j.isci.2022.103897](https://doi.org/10.1016/j.isci.2022.103897)

Provided by King's College London

Citation: Study finds potential predictor for perimenopause (2022, February 11) retrieved 23 May 2024 from <https://medicalxpress.com/news/2022-02-potential-predictor-perimenopause.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
