

# No time to exercise? What about three seconds a day?

February 7 2022

---



Credit: Pexels

Lifting weights for as little as three seconds a day can have a positive impact on muscle strength, a new study from Edith Cowan University (ECU) has discovered.

A collaboration with researchers from Niigata University of Health and Welfare (NUHW) in Japan had 39 healthy university students perform one [muscle contraction](#) at maximum effort for three seconds per day, for five days a week over four weeks.

The participants performed either an isometric, concentric or eccentric bicep curl (see definitions below) at maximum effort, while researchers measured the muscles' maximum voluntary contraction [strength](#) before and after the four-week period.

Another 13 students performed no exercise over the same period and were also measured before and after the four weeks.

Muscle strength increased more than 10 percent for the group who performed the eccentric bicep curl after the four weeks, but less increase in [muscle strength](#) was found for the other two exercise groups.

The no exercise group saw no increase.

Lead researcher Professor Ken Nosaka from ECU's School of Medical and Health Sciences said the results showed people didn't need to spend vast amounts of time exercising to improve their [muscle](#) strength.

"The study results suggest that a very small amount of exercise stimulus—even 60 seconds in four weeks—can increase muscle strength," he said.

"Many people think you have to spend a lot of time exercising, but it's not the case. Short, good quality exercise can still be good for your body and every muscle contraction counts."

## **Isometric vs concentric vs eccentric**

These three classifications relate to what the muscle is doing when being activated.

An isometric contraction is when the muscle is stationary under load, concentric is when the muscle is shortening and eccentric when the muscle is lengthening.

For a bicep curl, a dumbbell held with an arm by one's side, before lifting the weight upwards towards the chest and then lowering it back down via the elbow.

Lifting the weight sees the bicep in concentric contraction, lowering the weight sees it in eccentric contraction, while holding the weight parallel to the ground is isometric.

## **So which is best?**

The study shows all three lifting methods had some benefit to muscle strength, however eccentric contraction easily produced the best results.

Researchers measured each group's concentric, isometric and eccentric strength.

The concentric lifting group improved slightly (6.3 percent) in isometric strength but saw no improvement elsewhere, while the isometric group only saw an increase in eccentric strength (7.2 percent).

However, the eccentric group saw significant improvements in strength across all three measurements: concentric increased 12.8 percent, isometric 10.2 percent and eccentric 12.2 percent.

The eccentric group's overall muscle strength improved 11.5 percent after 60 seconds of effort in total.

"Although the mechanisms underpinning eccentric contraction's potent effects are not clear yet, the fact only a three-second maximal eccentric contraction a day improves muscle strength in a relatively short period is important for health and fitness," Professor Nosaka said.

## **Time-poor no more**

Professor Nosaka said the findings were exciting for promoting physical fitness and health, such as prevention of sarcopenia—a decrease in muscle mass and strength with aging.

"We haven't investigated other muscles yet, but if we find the three-second rule also applies to other muscles then you might be able to do a whole-body exercise in less than 30 seconds," he said.

"Also, performing only one maximal contraction per day means you don't get sore afterwards."

Professor Nosaka and NUHW's Dr. Masatoshi Nakamura designed the study and the data were collected by Dr. Nakamura and his Ph.D. and Masters students.

"Effect of daily 3-s maximum voluntary isometric, concentric or eccentric [contraction](#) on elbow flexor strength" was published in the *Scandinavian Journal of Medicine and Science in Sports*.

**More information:** Shigeru Sato et al, Effect of daily 3-s maximum voluntary isometric, concentric or eccentric contraction on elbow flexor strength, *Scandinavian Journal of Medicine & Science in Sports* (2022).

[DOI: 10.1111/sms.14138](https://doi.org/10.1111/sms.14138)

Provided by Edith Cowan University

Citation: No time to exercise? What about three seconds a day? (2022, February 7) retrieved 24 April 2024 from <https://medicalxpress.com/news/2022-02-seconds-day.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.