

The effect of dextromethorphan on cardiac Ca^{2+} handling and ion channels in a human iPSC model of TS. a, Representative traces of time-course Ca^{2+} imaging in spontaneously contracting cardiomyocytes from patients with TS treated with dextromethorphan (5 μM , until 120 min). b,c, Ca^{2+} transient frequency (b) and duration (c) analyses of cardiomyocytes from patients with TS before and after dextromethorphan treatment (n = 19). d, Representative traces of Ca^{2+} currents in cardiomyocytes from patients with TS with and without dextromethorphan. e, Late Ca^{2+} current analysis of cardiomyocytes from patients with TS with and without dextromethorphan treatment (TS, n = 11; with dextromethorphan, n = 12). f, Representative traces of Ba^{2+} currents in cardiomyocytes from patients with TS without treatment or treated with dextromethorphan (5 μM , 2 h, dextromethorphan) or with dextromethorphan and a SIGMAR1 antagonist, NE-100 (1 μM , dextromethorphan and NE-100). g, Voltage-dependent inactivation in cardiomyocytes from patients with TS without treatment (n = 25) or treated with dextromethorphan (n = 16) or with dextromethorphan and NE-100 (n = 11). h, Representative traces of IKr currents (E-4031 sensitive) in cardiomyocytes from patients with TS treated with dextromethorphan (5 μM) or dextromethorphan and NE-100 (each at 5 μM) or without treatment. i, IKr current amplitude analysis of cardiomyocytes from patients with TS treated with dextromethorphan (n = 9) or dextromethorphan and NE-100 (n = 10) or without treatment (n = 10) (*P

Citation: Cough suppressant knocks some hearts back into rhythm (2022, February 18) retrieved 17 May 2024 from <https://medicalxpress.com/news/2022-02-suppressant-hearts-rhythm.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.