


# Improving treatment for Parkinson's

February 18 2022

## Pedunculopontine nucleus (PPN)



Stimulation of glutamatergic neurons in the caudal PPN in the brainstem leads to recovery of locomotion in Parkinsonian mice. Credit: University of Copenhagen

Parkinson's is a neurodegenerative disease where dopaminergic neurons progressively die in the brainstem. Tremor and difficulties walking are recognizable movement symptoms for many people suffering from Parkinson's. Over time, nearly a quarter of patients will have so much trouble walking that they often end up freezing on the spot and falling, and many become housebound.

People are primarily treated with medicine, but in some cases doctors use Deep Brain Stimulation (DBS). In DBS, the surgeon places a thin metal wire in the brain, which can be used to send electrical pulses. DBS is effective in treating tremor, but alleviating difficulties in walking and freezing remains a challenge.

Now, a study from the University of Copenhagen conducted in mice demonstrate that DBS treatment of walking problems in Parkinson's could be optimized by targeting specific [neurons](#) in the brainstem—possibly benefitting some of the more than 7 to 10 million people suffering from the disease worldwide.

## **"Brainstem DBS are the right strategy to facilitate patients to walk properly again"**

Based on previous animal studies of motor circuits, which are responsible for the planning, control, and execution of voluntary movements, scientists have hypothesized that freezing of walking in Parkinson's could be alleviated. That would require DBS to stimulate neurons in the pedunclopontine nucleus (PPN), which is located in the brainstem. The PPN was thought to send signals from the brain to the spinal cord leading to body movements.

"However, initial results from [clinical trials](#) with DBS of the PPN had very variable effect on movement recovery, particularly in patients who

experience freezing of walking. It has therefore been debated where within the brainstem an optimal stimulation should be. Our study brings new knowledge to the table regarding the best area for DBS in order to alleviate this particular symptom," says corresponding author Professor Ole Kiehn at the Department of Neuroscience.

Previous results from the group showed that stimulation of so-called excitatory neurons in the PPN could initiate locomotion in normal mice. It raised the possibility that these nerve cells could indeed be used to treat movement symptoms in mice with features of Parkinson's Disease.

"We use a technology to target specific group of cells in the PPN in order to close in on what areas are the best to stimulate, if we want to alleviate these particular symptoms. The result shows that the motor improvement is optimal, if we stimulate what we call excitatory neurons in the caudal area of the PPN," explains Ole Kiehn.

"We believe that clinical trials with brainstem DBS are the right strategy to facilitate patients to walk properly again. But the variable clinical results occur, because DBS would require higher precision to target the particular group of neurons in the caudal PPN. It is a very delicate area, because if we were to stimulate excitatory neurons in other areas than the caudal PPN, it would cause complete immobilization instead."

## **The key is the activation of PPN neurons**

In Parkinson's Disease, [nerve cells](#) that produce dopamine progressively die. Since the 1960s, doctors have relied on medication to replace the missing dopamine, but it is notoriously difficult to fully control symptoms as the disease progresses.

"In many people the movement symptoms do not respond well to medical treatment in the later stages of this disease, so there has been

done a lot of research into alternative treatments, including a search for optimal targets for [deep brain stimulation](#)," explains Postdoc Debora Masini, first author of the new study, which included several different strategies to substantiate their findings.

"When we stimulated these specific neurons in the caudal area of the PPN, the animals were able to walk normally, across longer distances and with normal walking speed, as opposed to before the stimulation, where they would display symptoms of Parkinson's Disease," says Debora Masini.

"We systematically compared stimulation of different locations and cell types in a series of complementary experiments. And they all pointed towards the same conclusion. It strongly indicates these excitatory neurons in the caudal PPN are an ideal target for recovery of movement loss," she says.

The researchers hope that the new study could aid clinicians when they pick the exact location for DBS in the brainstem.

"The mice in our study only partially represents the complexity of this disease, but the results have been very telling. Nearly everything we have learned in the beginning on how to treat Parkinson's Disease comes from animal models, including the medication we use nowadays for patients. In this sense, it is a valid approach, and we hope our study can help provide better treatment for human patients," says Debora Masini.

The study was published in *Nature Communications*.

**More information:** Débora Masini et al, Targeted activation of midbrain neurons restores locomotor function in mouse models of parkinsonism, *Nature Communications* (2022). [DOI: 10.1038/s41467-022-28075-4](https://doi.org/10.1038/s41467-022-28075-4)

Provided by University of Copenhagen

Citation: Improving treatment for Parkinson's (2022, February 18) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-02-treatment-parkinson.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.