

Unraveling correlation of Cd induced metabolic toxicity and diabetes

February 21 2022



Representation of disturbed metabolic profiles by Cd exposure. Credit: Zhejiang University

Diabetes has become the leading metabolic disease threatening human health. Meanwhile, the association between exposure to harmful environmental factors and the occurrence of diabetes has always been a hot spot of public health and environmental medicine.



A recent study conducted by Dr. Zhou Zhou's lab at the Department of Environmental Medicine, the Zhejiang University School of Medicine elucidated the Cd perturbed metabolomic signature in <u>pancreatic beta</u> <u>cells</u> and human urine. The original article was published on line in *Environment International* on February 14, 2022.

Cadmium (Cd), as an extremely toxic heavy metal, exists widely in the environment. Cd pollution from industrial and agricultural activities is a worldwide environmental health threat to the public. The general population is exposed to Cd through foods such as rice, wheat, and potatoes from polluted agricultural soil. Tobacco smoking and seafood intaking further increase Cd exposure. Epidemiological studies indicated that urinary Cd content is markedly associated with impaired fasting glucose. However, there is still paucity of information on metabolic toxicity of Cd exposure in pancreatic β -cells. The underlying metabolic events of Cd exposure at metabolome-wide level in pancreatic β -cells and in humans remain largely unknown.

In this study, the researchers firstly delineated the metabolomic signatures of Cd exposure in insulin-secreting MIN6 cells and in urine of exposed workers by using non-target metabolomics. Then, they verified that these metabolic changes are linked to Cd toxicity and the perturbed metabolites and pathways correlates with pathogenesis of diabetes.

"Findings of this study greatly extend our understanding of metabolic toxicity of Cd exposure in pancreatic β -cells at metabolome-wide level and offer some new clues for linking Cd exposure to development of diabetes mellitus," as pointed out by Dr. Frederic Coulon, editor-in-chief of *Environment International*. Also, the results of this study support the notion that Cd induced metabolic toxicity could be monitored by examining perturbed urinary metabolites in humans and highlight the significance of reducing Cd exposure via drinking water at population level.



More information: Huihui Hong et al, Cadmium perturbed metabolomic signature in pancreatic beta cells correlates with disturbed metabolite profile in human urine, *Environment International* (2022). DOI: 10.1016/j.envint.2022.107139

Provided by Zhejiang University

Citation: Unraveling correlation of Cd induced metabolic toxicity and diabetes (2022, February 21) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2022-02-unraveling-cd-metabolic-toxicity-diabetes.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.