

Prescribing for anxiety has increased in those under age 35

March 22 2022



Credit: Pixabay/CC0 Public Domain

Researchers from the University of Bristol have found that there have been increases in incident prescribing of most anti-anxiety medications (called anxiolytics) in recent years, which have been substantial in young

adults (aged under 25).

This may reflect better detection of [anxiety](#), increasing severity of symptoms, increasing acceptability of [medication](#), or an earlier unmet need. While overall incident benzodiazepine prescribing has fallen over time, prescribing has increased in those aged under 35. In 2017, 44% of benzodiazepine prescriptions were longer than the NICE recommended maximum of four weeks.

Some of this prescribing is not based on robust evidence of effectiveness, such as the use of beta-blockers, some may contradict guidelines, such as anti-psychotics, and there is limited evidence on the effect of taking antidepressants long-term. As such, there may be unintended harm.

Prior to this research, previous studies had found substantial increases in the prescribing of antidepressants for any indication, and for depression, over the past two decades. However, little was known about trends in the prescribing of anxiolytics (antidepressants, benzodiazepines, beta-blockers, anticonvulsants, antipsychotics) for the treatment of anxiety. Several changes may have affected prescribing in recent years, including changes in clinical guidance.

Dr. Charlotte Archer, an NIHR-funded Senior Research Associate in Primary Care Mental Health at Bristol Medical School (PHS), said: "The increases in the number of new patients starting anxiolytic treatment may reflect better detection of anxiety by GPs. The increase in new prescriptions was substantial in those aged 18–34, particularly in recent years, and this may also reflect an earlier unmet need for young adults.

"It is also important to consider that some of this prescribing is not based on robust evidence of effectiveness and some may contradict guidelines. There is also limited evidence on the effect of taking antidepressants

long-term, so there may be unintended harm to those that are being prescribed this medication for a long time.

"The information from this study has allowed us to gain a better understanding of primary care prescribing for anxiety, and how trends have changed over the study period. Going forward, we need research to improve our understanding of why we are seeing this increase in prescribing, especially in [young adults](#), and to provide interventions that can reduce the growing reliance on medication for this age group."

More information: Charlotte Archer et al, Rise in prescribing for anxiety in UK primary care between 2003 and 2018: a population-based cohort study using Clinical Practice Research Datalink, *British Journal of General Practice* (2022). [DOI: 10.3399/BJGP.2021.0561](https://doi.org/10.3399/BJGP.2021.0561)

Provided by University of Bristol

Citation: Prescribing for anxiety has increased in those under age 35 (2022, March 22) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-03-anxiety-age.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
