

Study finds convalescent plasma can be effective early COVID-19 therapy

March 30 2022



A bag of antibody-rich COVID convalescent plasma after thawing. Credit: Will Kirk, Johns Hopkins University

The *New England Journal of Medicine (NEJM)* today published final results of a nationwide multicenter study led by researchers at Johns Hopkins Medicine and the Johns Hopkins Bloomberg School of Public Health that show plasma from patients who have recovered from COVID-19 and whose blood contains antibodies against SARS-CoV-2,

the causative virus, is an effective and safe option as an early outpatient treatment for the disease.

The research showed that high-titer (antibody-rich) COVID [convalescent plasma](#)—when administered to COVID-19 outpatients within nine days after testing positive—reduced the need for hospitalization for more than half of the study's predominantly unvaccinated outpatients. The U.S. Food and Drug Administration (FDA) currently authorizes this plasma as a [treatment option](#) for outpatients with immunocompromising diseases or receiving immunocompromising medications, and for all patients hospitalized with early-stage COVID-19.

The findings were first presented in a preprint [posted to MedRxiv](#) on Dec. 21, 2021.

"Based on our findings and conclusions—which are now validated through the peer-review process—we encourage [health care professionals](#) to keep SARS-CoV-2 antibody-rich blood plasma available in their blood banks as part of the [treatment](#) arsenal against early-stage COVID-19," says study co-lead author David Sullivan, M.D., professor of molecular microbiology and immunology at the Johns Hopkins Bloomberg School of Public Health with a joint appointment in [infectious diseases](#) at the Johns Hopkins University School of Medicine.

"We believe that the best role for convalescent plasma is extending its use to early outpatient treatment when other therapies, such as [monoclonal antibodies](#) or drugs, are either not readily available—as in low- and [middle-income countries](#)—or ineffective, as with SARS-CoV-2 variants that are resistant to certain monoclonal antibodies," Sullivan adds.

In the outpatient early-treatment study conducted between June 2020 and October 2021, the researchers provided 1,181 randomized patients

with one dose each of either polyclonal high-titer convalescent plasma (containing a concentrated mixture of antibodies specific to SARS-CoV-2) or placebo control plasma (with no SARS-CoV-2 antibodies). The patients were 18 and older, and had tested positive for SARS-CoV-2 within eight days prior to transfusion. A successful therapy was defined as a patient not requiring hospitalization within 28 days after plasma transfusion.

The study found that 17 patients out of 592 (2.9%) who received the convalescent plasma required hospitalization within 28 days of their transfusion, compared with 37 out of 589 (6.3%) who received placebo control plasma. This translated to a relative risk reduction for hospitalization of 54%.

Timing of the convalescent plasma transfusion also is critical: "The earlier the better," the researchers say.

"Based on the findings of an analysis in the new paper that wasn't available when the preprint was posted, we found that if convalescent plasma is given within five days after diagnosis, the effectiveness at reducing hospitalization approximated 80%," says Sullivan.

"We concluded that these results strongly support high-titer SARS-CoV-2 convalescent plasma as an effective early treatment for COVID-19 with advantages such as low cost, wide availability and rapid resilience to the virus's evolving variants," says study co-lead author Kelly Gebo, M.D., M.P.H., professor of medicine at the Johns Hopkins University School of Medicine.

The next step, the researchers say, is to make convalescent plasma for the outpatient treatment of COVID-19 easier to use, more efficiently administered and more accessible to those who might need it. As part of that effort, they have provided clinicians with a guide for implementing

a plasma transfusion center for outpatients with COVID-19, including logistical, staffing and blood banking requirements. [The guide appears in a paper](#) published March 29, 2022, in the journal *Transfusion*.

The team also continues to seek more understanding of what else convalescent plasma can do for outpatients with COVID-19. A soon-to-be published study will look at the ability of plasma to neutralize SARS-CoV-2 variants, including delta and omicron, despite no previous donor exposure to those viruses.

Since the study findings were first announced last December, there have been three developments supporting the use of convalescent plasma for early-stage COVID-19:

- On Dec. 28, 2021, [the FDA expanded](#) the authorized emergency use of convalescent plasma with high titers of anti-SARS-CoV-2 antibodies "for the treatment of COVID-19 in patients with immunosuppressive disease or receiving immunosuppressive treatment, in either the outpatient or inpatient setting."
- On Feb. 2, 2022, [the Infectious Disease Society of America updated](#) its "Guidelines on the Treatment and Management of Patients with COVID-19" to include the "use of convalescent plasma in ambulatory patients with mild-to-moderate COVID-19 at high risk for progression to severe disease with no other treatment options."
- On March 7, 2022, [the American Red Cross announced](#) that it was "temporarily testing all blood donations for COVID-19 antibodies to help identify donations that could be processed into convalescent plasma." The organization said this was being done "to help support immunocompromised patients battling COVID-19."

"These recent acknowledgements of high-titer convalescent [plasma](#)'s benefit in treating early-stage COVID-19—in conjunction with our peer-reviewed findings and our new guide for more effective administration of the treatment—provide clinicians with an additional option for outpatients," Gebo says.

More information: Early Outpatient Treatment for Covid-19 with Convalescent Plasma, *New England Journal of Medicine* (2022). www.nejm.org/doi/full/10.1056/NEJMoa2119657

Evan M. Bloch et al, How do I implement an outpatient program for the administration of convalescent plasma for COVID -19?, *Transfusion* (2022). [DOI: 10.1111/trf.16871](https://doi.org/10.1111/trf.16871)

Provided by Johns Hopkins University School of Medicine

Citation: Study finds convalescent plasma can be effective early COVID-19 therapy (2022, March 30) retrieved 27 April 2024 from <https://medicalxpress.com/news/2022-03-convalescent-plasma-effective-early-covid-.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.