

# Origins of diabetes may be different in men and women, according to new research

March 23 2022

---



Credit: Pixabay/CC0 Public Domain

Over the past four decades, global cases of Type 2 diabetes mellitus have skyrocketed. According to the World Health Organization, the number of people estimated to have the disease jumped from 108 million in

1980 to 422 million in 2014, with the fastest growth observed in low- and middle-income countries.

Although the disease is common, there is still much research left to be done to fully understand it. For instance, while diabetes is linked to obesity, researchers still do not know the exact reasons why obesity causes diabetes.

In a new paper published in the journal *Obesity Reviews*, Concordia researchers Kerri Delaney and Sylvia Santosa look at how fat tissue from different parts of the body may lead to diabetes onset in men and women. They reviewed almost 200 hundred [scientific papers](#) looking for a deeper understanding of how fat operates at the surface and tissue level, and the mechanisms by which that tissue contributes to diabetes onset.

"There are many different theories about how diabetes develops, and the one that we explore posits that different regions of fat tissue contributes to [disease risk](#) differently," says Kerri Delaney, a Ph.D. candidate at Concordia's PERFORM Centre and the paper's lead author. "So the big question is, how do the different depots uniquely contribute to its development, and is this contribution different in men and women?"

## **From surface to cell level**

Men and women store fat in [different places](#). Diabetes, like many other diseases, is closely associated with abdominal fat. Women tend to store that fat just under the skin. This is known as subcutaneous fat. In men, [abdominal fat](#) is stored around the organs. This is visceral fat.

Fat appears to exhibit different features in men and women. They grow differently, are dispersed differently and interact with the inflammatory and immune system differently. For example, in men [fat tissue](#) expands

because the [fat cells](#) grow in size; in women, fat cells multiply and increase in number. This changes with the loss of the protective hormone estrogen that disappears with menopause and may explain why men are more susceptible to diabetes earlier in life than women.

Working from the hypothesis that diabetes risk is driven by expansions of visceral fat in men and of subcutaneous fat in women, the researchers then looked through the papers to see what was happening in the cell-level microenvironments.

Though more research is needed, there were overall differences observed in the immune cell, hormone, and cell signaling level in men and women that seem to support different origins in diabetes between the sexes.

Delaney and Santosa hope that by identifying how diabetes risks are different in men and women, clinical approaches to treatment of the disease can be better defined between the sexes.

"Currently, the treatment of diabetes is similar for men and women," says Santosa, an associate professor in the department of Health, Kinesiology and Applied Physiology. "If we understood the differences between them better, we could consider these mechanisms in recommending treatments to men and [women](#) based on how diabetes medications work."

**More information:** Kerri Z. Delaney et al, Sex differences in regional adipose tissue depots pose different threats for the development of Type 2 diabetes in males and females, *Obesity Reviews* (2022). [DOI: 10.1111/obr.13393](https://doi.org/10.1111/obr.13393)

Provided by Concordia University

Citation: Origins of diabetes may be different in men and women, according to new research (2022, March 23) retrieved 25 May 2024 from <https://medicalxpress.com/news/2022-03-diabetes-men-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.