

# First human study using sound waves to break up kidney stones shows promising results

March 22 2022

---



Credit: Unsplash/CC0 Public Domain

An innovative technique called burst wave lithotripsy (BWL) may provide an effective, more accessible alternative for noninvasive treatment of kidney stones, according to initial human studies reported in the *Journal of Urology*, an Official Journal of the American

Urological Association (AUA).

Using focused ultrasound bursts to break up [kidney stones](#), BWL provides a promising new approach to avoiding the [high costs](#) and healthcare burden of kidney [stone](#) treatment, according to the report by Jonathan Harper, MD, of University of Washington School of Medicine and colleagues. They write, "New BWL technology successfully fragmented stones of a variety of sizes, locations and densities to under two-millimeter fragments within 10 minutes with negligible tissue injury."

## **Toward new office-based treatment approach for kidney stones**

Kidney stones are a very common condition, affecting 1 in 10 Americans at a cost of \$10 billion per year. While many stones pass on their own, other treatments are sometimes needed. Some fairly small stones can be treated using a technique called extracorporeal shock wave lithotripsy (ESWL), using shock waves to break up the stones so that they will be more likely to pass on their own. The ESWL procedure is performed in a hospital or clinic, typically with the patient under sedation.

In contrast to the [shock waves](#) used in ESWL, the BWL procedure uses "short harmonic bursts" of ultrasound energy—potentially allowing the stones to be broken up in a shorter procedure without the need for sedation or anesthesia. Pre-clinical studies supported the effectiveness of BWL in breaking up experimental stones of varying size and composition.

For the next step in developing the procedure, Dr. Harper and colleagues performed initial studies in [human patients](#) with kidney stones. The

patients were undergoing a surgical procedure, called ureteroscopy, that is used to treat larger stones. Before that treatment, the stones were treated with a short period of BWL—no longer than 10 minutes. Using the ureteroscope, the researchers were able to directly observe how well the ultrasound waves worked in breaking up stones, as well as observing any injury to the kidney tissues.

In the study, BWL was used in 19 patients to target 25 confirmed stones. Overall, a median of 90% of the stone volume was completely broken up (comminuted) within the brief treatment period. Thirty-nine percent of stones were completely fragmented while 52% were partially fragmented.

Most of the stone fragments measured less than two millimeters—small enough that they would be easily able to pass on their own, without much pain. Examination through the ureteroscope showed mild or no tissue injury related to the BWL procedure, consisting mainly of mild bleeding.

For several reasons, BWL could be an important practical advance in the management of kidney stones. Many patients make multiple visits to the emergency department while waiting for definitive treatment of their kidney stones, often requiring opioids for pain management. Some patients require repeated treatment, even after surgery or ESWL. "The ability to noninvasively break stones and expel the fragments in awake patients at first presentation in the ED or clinic has the potential to provide just-in-time treatment leading to a reduction of the overall pain, cost and resource burden associated with a stone event," researchers write.

The new results "are a step toward an office-based lithotripsy for awake patients," Dr. Harper and coauthors conclude. They plan further studies toward determining whether the BWL technique can meet their ultimate

goal of a "noninvasive, 30-minute treatment in the clinic without anesthesia."

**More information:** Jonathan D. Harper et al, Fragmentation of Stones by Burst Wave Lithotripsy in the First 19 Humans, *Journal of Urology* (2022). [DOI: 10.1097/JU.0000000000002446](https://doi.org/10.1097/JU.0000000000002446)

Provided by Wolters Kluwer Health

Citation: First human study using sound waves to break up kidney stones shows promising results (2022, March 22) retrieved 25 April 2024 from <https://medicalxpress.com/news/2022-03-human-kidney-stones-results.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.