


## Inflammation and net-like protein structures in cerebral cavernous malformations

## March 30 2022



Proinflammatory cytokines and chemokines mainly associated with recruitment of neutrophils and monocytes were elevated in the cerebellum of Ccm3<sup>iECKO</sup> mice (CCM3) compared with wild-type controls (WT) during acute CCM. The following cytokines and chemokines are shown: IL-1 $\beta$  (A), IL-6 (B), TNF (C),



CXCL1/KC/GRO (D), CXCL2/MIP-2 (E), CCL2/MCP-1 (F), CCL3/MIP-1 $\alpha$  (G) and CXCL10/IP-10 (H), at P6 (n = 4–5 per group) and P8 (n = 5–9 per group). I Representative images of the whole brain at P6 and P8. \*P

Citation: Inflammation and net-like protein structures in cerebral cavernous malformations (2022, March 30) retrieved 9 April 2024 from <a href="https://medicalxpress.com/news/2022-03-inflammation-net-like-protein-cerebral-cavernous.html">https://medicalxpress.com/news/2022-03-inflammation-net-like-protein-cerebral-cavernous.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.