


# Largest genetic study of pleural infection reveals complex nature of common disease

March 15 2022

---



Credit: CC0 Public Domain

A new genetic study of the bacteria that cause pleurisy has shown most cases involve more than one type of bacterium, and revealed which bacterial combinations cause the most serious infections.

Pleural infection is a common and severe disease with high morbidity and mortality worldwide. Pleural infection occurs when bacteria invade the pleural space, which is the area in-between the lungs and the chest wall. Patients need hospitalization, which could last from several days to weeks. Conventional bacterial culture techniques fail to identify the pathogen in approximately 40% of cases, which leads to broad-spectrum antibiotics being used as treatment.

In a new study published in *The Lancet Microbe*, an international collaboration of scientists led by researchers from Oxford University's Nuffield Department of Medicine has found that pleural infection was predominately caused by more than one microbe at a time.

The researchers used genetic analysis to study the entire range of bacteria in patients with pleural infection, to help them assess the correlation between bacterial patterns and 1-year patient survival. This is the largest translational metagenomics study of pleural infection in adults, combining high-throughput discovery with high-quality prospective clinical data. The study used samples collected from a previous clinical trial led by John Corcoran and Professor Najib Rahman.

The study was jointly led by Najib Rahman, Professor of Respiratory Medicine, and Nikolaos Kanellakis of the Nuffield Department of Medicine's Experimental Medicine Division.

Nikolaos Kanellakis says that "improving our understanding of the underlying biology of pleural infection has the potential to improve patients' clinical management and potentially shorten hospital stays, minimize complications from antibiotic use, and reduce health-care costs.

"Understanding the crosstalk between human and bacterial cells, and the

interactions between bacteria to form [biofilms](#), might also help us to develop non-antibiotic-based treatments in the future.

"Importantly, the establishment and use of a method for identifying [pathogens](#) without using the slower method of culturing bacterial plates in the lab, could one day lead to patients being given the most appropriate treatment much faster."

The study also showed mixed anaerobes and other Gram-negative bacteria dominating community-acquired polymicrobial infection, whereas *Streptococcus pneumoniae* dominated in monomicrobial cases. Infections with anaerobes and bacteria of the *Streptococcus anginosus* group were associated with better survival, whereas *Staphylococcus aureus* and *Enterobacteriaceae* were linked with higher mortality.

**More information:** Nikolaos I Kanellakis et al, The bacteriology of pleural infection (TORPIDS): an exploratory metagenomics analysis through next generation sequencing, *The Lancet Microbe* (2022). [DOI: 10.1016/S2666-5247\(21\)00327-X](https://doi.org/10.1016/S2666-5247(21)00327-X)

Provided by University of Oxford

Citation: Largest genetic study of pleural infection reveals complex nature of common disease (2022, March 15) retrieved 23 April 2024 from <https://medicalxpress.com/news/2022-03-largest-genetic-pleural-infection-reveals.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.