

Pain neuron-derived peptide prevents endotoxic death by targeting kynurenine pathway in microglia

March 8 2022





Pain neuron-derived Reg3γ protects brain metabolism of LPS challenged mice. Credit: Kenta Maruyama

A research team led by the National Institute for Physiological Sciences and joined by Hokkaido University explored the role of pain neurons in the regulation of endotoxic death. They found that peptide named Reg 3γ acts as a pain neuron-enriched brain-targeted hormone that protects the host from endotoxic death.

Sepsis is a high-mortality disease that arises when the body's immune reaction to pathogens causes multi-organ defects. Although infectioninduced pro-inflammatory cytokines are indispensable for pathogen elimination, dysregulated production of these factors can lead to endotoxic shock. Despite the extensive use of anti-TNF- α antibody administration or glucocorticoids in patients undergoing endotoxic shock, mortality rates remain high at 30%. These disappointing results suggest that the mechanism of endotoxic death is only partially explained by uncontrolled inflammation.

Studies of <u>pain</u> neurons have traditionally focused on the <u>ion channels</u>. Natural pain sensors such as the transient receptor potential cation channel subfamily V member 1 (TRPV1), and TRP ankyrin 1 (TRPA1) are expressed in afferent pain neurons and trigger pain to protect the organism from further harm. Emerging reports suggest that organinnervated pain neurons and TRP channels also regulate dermal innate immunity, psoriasis, islet function, candidiasis, and osteomyelitis. Mechanistically, these phenomena are thought to be evoked by the paracrine secretion of nociceptor-derived peptides such as Calcitonin gene-related peptide (CGRP) and VIP, which modulate immune signals and endocrine pathways. These findings have also raised the question of



whether pain neuron-derived "hormones" exist.

In this new study, Dr. Kenta Maruyama at NIPS, Assistant Professor Takeshi Kondo at Hokkaido University, and colleagues revealed that pain neuron-derived peptide Reg3 γ penetrates the inflamed brain and suppresses the expression of microglial IDO1, a key enzyme of the kynurenine pathway. Endotoxin-administered pain neuron-null mice and pain neuron-specific Reg3 γ deficient mice exhibit a high-mortality rate accompanied by decreased brain HK1 phosphorylation and ATP production despite normal inflammation. This metabolic arrest is only observed in the brain, and aberrant production of brain quinolinic acid, a neurotoxic metabolite of the kynurenine pathway, causes HK1 suppression. Notably, brain administration of Reg3 γ protects mice from endotoxic death by enhancing <u>brain</u> ATP production. By identifying pain neuron-derived Reg3 γ as a microglia-targeted hormone, this discovery provides novel insights into the understanding of tolerance to endotoxic death.

The study is published in *Cell Reports*.

More information: Kenta Maruyama, Nociceptor-derived Reg3g prevents endotoxic death by targeting kynurenine pathway in microglia, *Cell Reports* (2022). DOI: 10.1016/j.celrep.2022.110462

Provided by National Institutes of Natural Sciences

Citation: Pain neuron-derived peptide prevents endotoxic death by targeting kynurenine pathway in microglia (2022, March 8) retrieved 3 May 2024 from https://medicalxpress.com/news/2022-03-pain-neuron-derived-peptide-endotoxic-death.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.