

Army Guard soldiers, especially those with combat exposure, are at risk for alcohol misuse after military deployment

March 9 2022, by Kimberly Flynn



Combat exposure can increase the risk for alcohol misuse after military deployment. Credit: Rutgers Center of Alcohol & Substance Use Studies

Using alcohol to cope with negative emotions after military deployment is known to be common among active-duty servicemembers. But a new study in the *Journal of Studies on Alcohol and Drugs* suggests that Army

Guard soldiers who return home after deployment may face similar risks.

Further, the situation might be more troublesome for Guard soldiers who were exposed to combat.

"Combat events, such as having engaged in direct combat and experienced combat trauma, may precipitate a great deal of personal discomfort ('moral injury'), necessitating some form of self-soothing, such as excessive alcohol use," says study author James Griffith, Ph.D., a faculty member and research fellow at the University of Utah's National Center for Veterans Studies.

In his research, Griffith looked at survey responses from Army National Guard soldiers who had recently returned from Operation Iraqi Freedom in 2010.

Griffith used data from more than 4,500 Army National Guard soldiers who responded to the U.S. Army's Reintegration Unit Risk Inventory. The anonymous survey includes 80 questions about alcohol and drug use, as well as other aspects of the soldiers' lives such as criminal behavior, interpersonal relationships, social support and psychological distress symptoms.

He compared their responses to those of Guard soldiers serving their traditional part-time military role, as well as to previous research on deployed Army Guard and active-duty military personnel.

Overall, recently deployed and returned Army Guard soldiers had higher rates of heavy drinking than nondeployed Army Guard soldiers serving in a traditional, garrison role (29.9% vs. 24.1% for heavy drinking, and 33.9% vs. 31.8% for binge drinking, respectively). The rate of drinking among deployed Guard soldiers also exceeded that for deployed military

personnel in several past studies.

The most impactful factor was exposure to combat during deployment, which was associated with greater alcohol use.

Griffith says that the nature of the reserve military service may help contribute to its members being at higher risk for alcohol use. Most of the time, Army Guard personnel serve one weekend a month and 15 annual active-duty training days, as do most reservists. When called up, they serve full-time for a period of about one year. And in the recent Iraq and Afghanistan wars, Army Guard and Army Reserve personnel constituted about 30% to 40% of the U.S. force.

"On their return, Guard personnel resume their part-time military service and civilian life and employment," he says. "Unlike active-duty military personnel, Guard personnel typically do not live near military installations to receive behavioral health care. And many are not eligible for military health care unless conditions are directly related to active-duty military service."

Griffith says more needs to be done to help recently returning, deployed Army Guard soldiers, such as more thoroughly assessing alcohol risk. He also points out that the risk of unhealthy alcohol use is greatest within one year of returning—suggesting that a postdeployment educational program for servicemembers and their families on the risk factors for alcohol use could be helpful.

"Content would familiarize participants with indicators of unhealthy alcohol use and possible follow-up actions for those at risk," he says. "Specifically, information and guidance for obtaining informal and formal behavioral health care might be provided when soldiers are initially screened risk positive for alcohol misuse."

Special care needs to be taken to assess the emotional state of soldiers who have had traumatic combat experience, says Griffith, since they are at especially high risk of alcohol misuse.

"Given that negative emotions likely play a role in the connection of combat events and alcohol use, some form of screening should assess soldiers' emotional states, and for more severe negative emotions, include referral to psychological health care," Griffith says.

More information: Griffith, J. (2022). Postdeployment alcohol use and risk associated with deployment experiences, combat exposure, and postdeployment negative emotions among Army National Guard Soldiers. *Journal of Studies on Alcohol and Drugs*, 83, 202–211. [DOI: 10.15288/jsad.2022.83.202](https://doi.org/10.15288/jsad.2022.83.202)

Provided by Journal of Studies on Alcohol and Drugs

Citation: Army Guard soldiers, especially those with combat exposure, are at risk for alcohol misuse after military deployment (2022, March 9) retrieved 11 May 2024 from <https://medicalxpress.com/news/2022-03-reservists-combat-exposure-alcohol-misuse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.