

How much is too much for assistance level using robotic therapy in stroke rehabilitation?

March 25 2022



The stroke patient performing upper limb robotic therapy with tailor-made assistance from a robotic device with feedback displayed on the screen. Credit: Takashi Takebayashi , Osaka Prefecture University


The Department of Occupational Therapy, College of Health and Human Sciences, Osaka Prefecture University, Japan conducted exploratory research to examine the impact of robotic assistance in the upper limb rehabilitation of stroke survivors. Upper limb motor recovery is always a big challenge for rehabilitation professionals and also for the patient. In the past, there have been numerous theories proposed by many pioneers in stroke management; however, many of these theories failed. Currently, functional rehabilitation has been proven to be most effective in upper limb rehabilitation as it emphasizes meaningful task-specific training and practice-based learning.

Advances in technology like robotics have made therapists' work easier by assisting patients in doing guided limb movements and practicing the same with more repetition. However, there are contradictory opinions regarding the level of assistance that is ideal for a better prognosis. Researcher Takashi Takebayashi says that whether "robotic therapy can produce better upper extremity recovery compared to conventional rehabilitation remains unclear."

To answer this controversy a retrospective sub-analysis of a previous study was conducted. Takashi Takebayashi and colleagues selected an experimental study performed with 60 participants who were selected from 715 [stroke](#) patients of a wide age range using specific selection criteria, from six [rehabilitation](#) hospitals in Japan, during November 2008 to April 2010. The researchers randomly allocated the selected

patients into two groups, one receiving maximally guided [robotic therapy](#) and the other self-guided minimally assisted robotic exercises. In addition to collecting a detailed history of individual patients, performance-based impairment; upper limb motor functions and temporal performance of upper limb were also measured using standard outcome measures before and after 6 weeks of intervention.

Past research has suggested that maximal assistance using robotics may hamper the voluntary efforts generated by the patients and hence the assistance level should be within a tailor-made limit. As this claim is also contradicted by a few researchers, Takashi Takebayashi and colleagues wanted to solve the puzzle. So, they categorized the patients into high robotic assistance and little robotic assistance, based on the amount of external support and guidance offered by the robot.



The amount of assistance by the robot should be increased for subjects with severe-to moderate UE impairment, and the amount of assistance by the robot should be reduced for subjects with moderate-to-mild UE impairment. Credit: Takashi Takebayashi, Osaka Prefecture University

Both the groups underwent intervention for a 6-week period, and for avoiding bias a specialized person, who was blinded to the study, was used to conduct all the outcome parameters. The subjects were divided into many clusters among which the severity of stroke manifestation was the predominant one. The subjects were divided into mild to moderate groups and moderate to severe groups to understand whether there was a difference in the contribution of robotics among them.

The findings in their study demonstrated that the intervention group comprising stroke survivors with severe-to-moderate stroke severity (FMA control group showed significantly better improvement using the same outcome tools compared to the intervention group.

The researchers quoted from the study results that "to maximize the improvements of the affected UE function in sub-acute stroke survivors, the robotic assistance may need to be increased in patients with severe-to-moderate paralysis and decreased in patients with moderate-to-mild paralysis, to improve the performance of voluntary movements as much as possible." Hence, the research highlighted that selecting the level of intensity and assistance level offered by robotics is highly technical and professional, and hence using robotics needs more expertise and patient assessment. The researchers also bring to light the varied results produced by the end-effector and exoskeleton-type robotics, however, the researcher employed the end-effector type of robotics for his research and have produced this novel result.

The researchers draw support from previous studies that have stated that robotics can enhance the activation of the sensorimotor cortex during the intervention period and thereby improve motor performance significantly. The authors hypothesize that the changes that have happened among the participants might be because of the Hebbian plasticity. Having said this the researcher also advises future results to concentrate on proving the same using more objective tools like [functional magnetic resonance](#) imaging using larger samples.

To summarize, the researchers have clearly stated that "an optimal amount of robotic assistance was found to be a key to maximize improvement in post-stroke UE paralysis." The study is published in *Journal of NeuroEngineering and Rehabilitation*.

More information: Takashi Takebayashi et al, Impact of the robotic-assistance level on upper extremity function in stroke patients receiving adjunct robotic rehabilitation: sub-analysis of a randomized clinical trial, *Journal of NeuroEngineering and Rehabilitation* (2022). [DOI: 10.1186/s12984-022-00986-9](#)

Provided by Osaka Prefecture University

Citation: How much is too much for assistance level using robotic therapy in stroke rehabilitation? (2022, March 25) retrieved 2 May 2024 from <https://medicalxpress.com/news/2022-03-robotic-therapy.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--