


Study supports shortened quarantine period among asymptomatic health care professionals exposed to SARS-CoV-2

March 24 2022



Creative rendition of SARS-CoV-2 particles (not to scale). Credit: National Institute of Allergy and Infectious Diseases, NIH

New data published today in the *American Journal of Infection Control* (AJIC), suggest that healthcare professionals (HCPs) exposed to SARS-

CoV-2 can safely return to work after seven days of quarantine if they are asymptomatic and receive a negative COVID-19 test. The findings, from the first reported study evaluating reduced quarantine duration in a healthcare setting, offer a potential new strategy for mitigating staffing shortages prior to a next wave of COVID-19 cases.

"Maintaining appropriate staffing in [healthcare facilities](#) is essential to increasing safety for both HCPs and patients," said Graham M. Snyder, MD, MS, University of Pittsburgh School of Medicine and UPMC Presbyterian, and senior author on the published study. "The COVID-19 pandemic will likely persist for years, and the findings from our study suggest that a thoughtfully designed program can enable certain HCPs to return to work seven days following a higher-risk SARS-CoV-2 exposure, alleviating the serious workforce shortages we experienced over the past two years."

Current CDC guidelines recommend self-isolation at home for 14 days following a higher-risk SARS-CoV-2 exposure to reduce the potential for transmission. However, the risk of developing symptomatic COVID-19 after seven days post-exposure is low, and CDC guidelines also suggest alternate approaches when workforce shortages exist. In January 2021, UPMC changed their HCP SARS-CoV-2 quarantine policy from 14 to 10 days, based on internal data demonstrating an extremely low rate of COVID-19 symptom development after 10 days post-exposure.

To evaluate the potential for asymptomatic HCPs to safely return to work (RTW) after a shorter quarantine period, Dr. Snyder and colleagues designed and implemented an evidence-based program in which HCPs returned to work eight days after exposure if they remained asymptomatic through day seven and if they had a negative result from a SARS-CoV-2 nucleic acid amplification test (NAAT) taken five to seven days post-exposure. The researchers then evaluated this program

as a strategy for reducing workforce shortages during the COVID-19 pandemic.

Between January and June 2021, the study enrolled 384 unvaccinated HCPs who had a higher-risk workplace or non-household community SARS-CoV-2 exposure, were asymptomatic, and were

Citation: Study supports shortened quarantine period among asymptomatic health care professionals exposed to SARS-CoV-2 (2022, March 24) retrieved 27 April 2024 from <https://medicalxpress.com/news/2022-03-shortened-quarantine-period-asymptomatic-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.