

Smoking in underserved communities nearly double the US national rate

March 7 2022



Credit: CC0 Public Domain

Although rates of smoking and smoking-related cancers have declined in the United States in recent years, some people may be at higher risk for cigarette use and nicotine addiction and could therefore gain significant

benefits from tobacco prevention and cessation programs. New research indicates that the prevalence of smoking in underserved communities is nearly double the national prevalence, and that smoking is linked with mental health conditions and substance use disorders in these communities. The findings are published by Wiley online in *Cancer*.

For the study, Sue C. Lin, Ph.D., MS, of the Health Resources and Services Administration within the U.S. Department of Health and Human Services, analyzed results from the 2014 Health Center Patient Survey to assess the prevalence of smoking among adults from underserved communities who received primary care at federally qualified health centers. These centers serve individuals and families from underserved communities including people experiencing homelessness, agricultural workers, and residents of public housing. The team also examined associations of smoking with co-occurring mental health conditions and substance use disorders.

Among the major findings:

- The prevalence of smoking among adults from underserved communities was 28.1%, compared with 14.0% reported in the general U.S. population.
- Among those who currently smoked, 59.1% had depression and 45.4% had anxiety.
- Non-Hispanic Black adults who smoked had more than two times the odds of reporting substance use disorders.
- Individuals at or below 100% of the [federal poverty level](#) had more than two times the odds of having mental health conditions, and those who were unemployed had more than three times the odds for substance use disorders.

"Our study underscores the importance of understanding the association and increased risk of [mental health conditions](#) and [substance use](#)

[disorders](#) among adults from underserved communities who smoke while also addressing socioeconomic risk factors to achieve better health outcomes," said Dr. Lin. "The study further highlights the significance of tailored smoking cessation treatments for individuals from underserved communities that will support cancer prevention care."

More information: Sue C. Lin et al, Disparities in smoking prevalence and associations with mental health and substance use disorders in underserved communities across the United States, *Cancer* (2022). [DOI: 10.1002/cncr.34132](#)

Provided by Wiley

Citation: Smoking in underserved communities nearly double the US national rate (2022, March 7) retrieved 26 June 2024 from <https://medicalxpress.com/news/2022-03-underserved-national.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.