

Taking care of the young caregivers

March 15 2022



During the pandemic, many adolescent carers experienced growing pressures and loss of a break from their caring role. Credit: Tohoku University

Adolescent caregivers in the UK were more at risk of experiencing psychological distress and poor mental well-being during the COVID-19 pandemic than their non-caregiver peers. The finding, by Tohoku University researchers in Japan, was published in the *Journal of*

Adolescent Health and highlights the urgent need for better support systems for young caregivers.

There are approximately 800,000 young caregivers in the UK, defined as anyone under the age of 18 who looks after someone who is ill, disabled or elderly in need of care, without being paid. During the COVID-19 pandemic, most adolescent caregivers in the UK experienced growing pressures due to school closures, the disruption of established routines and a lack of breaks from [home life](#) and caring roles.

"While some studies have looked at the short-term effects of the first UK national lockdown, little is known about the long-term consequences of COVID-19 on [adolescent health](#)," says Miharu Nakanishi at the Tohoku University Graduate School of Medicine.

Nakanishi and colleagues in Japan and the UK examined the connection between the caring status of adolescents and their [mental health](#) outcomes before and during the pandemic.

They analyzed data from 3,927 people who took part in two surveys, including 3,616 non-caregivers and 311 adolescent caregivers. The Millennium Cohort Study is collecting data on the physical and mental health of 18,818 people born across the UK between 2000 and 2002. The COVID-19 survey was launched in May 2020 to explore the pandemic's effect on the lives of people from different generations and backgrounds.

The researchers combined data from the two surveys to explore how the [psychological symptoms](#) and mental well-being of young caregivers were impacted by the pandemic. Psychological symptoms were measured using the Kessler Distress Scale, which asks participants to rate their own anxiety and depressive symptoms. Mental well-being was measured using the Warwick-Edinburgh Mental Well-being Scale, a set of

questions designed to test different aspects of positive mental health.

They found that, compared with non-caregivers, the adolescent caregivers in the study reported higher [psychological distress](#) and lower mental well-being scores during the pandemic. Compared with non-caregivers, adolescent caregivers experienced significantly lower scores for [social support](#) and higher scores for loneliness

"Our study highlights the urgent need for better psychosocial support for adolescent caregivers," says Nakanishi. "They need systems in place to help them return to school and catch up with their studies and other students, such as psychoeducational sessions to build psychological resilience."

Next, the research team plans to collect data on the [mental health outcomes](#) of young caregivers in the Tokyo Teen Cohort, Japan during the pandemic. Comparing the results between countries will reveal more information about how different levels of COVID-19 restrictions impact mental health in early adulthood.

More information: Miharu Nakanishi et al, Adolescent Carers' Psychological Symptoms and Mental Well-being During the COVID-19 Pandemic: Longitudinal Study Using Data From the UK Millennium Cohort Study, *Journal of Adolescent Health* (2022). [DOI: 10.1016/j.jadohealth.2022.01.228](#)

Provided by Tohoku University

Citation: Taking care of the young caregivers (2022, March 15) retrieved 25 April 2024 from <https://medicalxpress.com/news/2022-03-young-caregivers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.