

Adults with asthma have a higher risk of developing obesity

April 27 2022



Credit: Unsplash/CC0 Public Domain

A study with more than 8,700 participants from 11 European countries and Australia has concluded that adults with asthma are more likely to become obese later in life. The research, led by the Barcelona Institute

of Global Health (ISGlobal), a center supported by the "la Caixa" Foundation, has also shown the risk is higher among patients with non-allergic asthma, those who have had the disease for a longer period or those who are under treatment with corticosteroids. These results have been published in *Thorax*.

"Several studies have shown that asthma and [obesity](#) share some common socioeconomic, behavioral and environmental risk factors that can lead to the development of both diseases. Some previous research focused on the mechanisms by which obesity could lead to asthma, but the inverse relationship had not received much attention until recently," says Subhabrata Moitra, who carried out this research at ISGlobal and is now a researcher at the University of Alberta.

The researchers used data from the cohort study European Community Respiratory Health Survey (ECRHS), collected in three waves between 1990 and 2014 and with follow-up visits at intervals of approximately 10 years. The different waves gathered data from questionnaires, lung function tests and measures taken to determine participants' body mass index and asthma status and characteristics.

Between the first and the second follow-ups, 14.6% of non-asthmatic participants developed obesity, while the percentage increased up to 16.9% among asthmatic participants. The difference between the two groups became more obvious when the research team translated it into a relative risk that took into consideration the role of other factors such as asthma and smoking, and saw that asthmatic had an increased 21% risk of obesity compared to non-asthmatic participants.

Asthmatic participants with longer [disease duration](#) had 32% larger risk of obesity than those with shorter duration and those with non-allergic asthma had an increased 47% risk than those with allergic asthma. Those participants treating the respiratory disease with corticosteroids showed a

99% higher risk of obesity compared to those who did not use this treatment, which had already been associated with abnormal weight gain in previous research.

"A potential explanation for the weight gain associated with asthma could be the reduction of physical activity in asthmatic patients. However, our results do not support this hypothesis, since the levels of physical activity in our study did not affect the observed association," says Judith Garcia-Aymerich, Head of the Non-Communicable Diseases and Environment Programme at ISGlobal and senior author of the study. "Regardless of the mechanisms, still unknown, our results have implications for the clinical care of adults with asthma," she adds.

Unlike a previous study that found this association between asthma and weight gain only in women, in this case there were no sex differences. Another previous study had found an association between [asthma](#) in children and obesity after a 10-year follow-up, but this is the first known study to show a similar association in adults regardless of their sex.

More information: Long-term effect of asthma on the development of obesity among adults: an international cohort study, ECRHS, *Thorax* (2022). [DOI: 10.1136/thoraxjnl-2021-217867](https://doi.org/10.1136/thoraxjnl-2021-217867)

Provided by Barcelona Institute for Global Health

Citation: Adults with asthma have a higher risk of developing obesity (2022, April 27) retrieved 7 May 2024 from <https://medicalxpress.com/news/2022-04-adults-asthma-higher-obesity.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--