

## Researchers study how to unlock clinical riskprediction models so they can be applied to multiple clinical settings

April 12 2022



Credit: Chokniti Khongchum from Pexels

Modern predictive models require large amounts of data for training and evaluation, the absence of which may result in models that are specific to



certain locations, their populations and the clinical practices there. Currently, best practices for clinical risk prediction models lack a level of "generalizability" that could vastly increase their usefulness for other clinical settings in other locations.

A team of NYU Tandon researchers led by Rumi Chunara, professor of computer science and engineering with affiliations with the NYU School of Global Public Health, investigated whether mortality prediction models vary significantly when applied to hospitals or geographies different from the ones in which they are developed. They also queried the data to determine specific characteristics of the datasets—involving analysis of electronic health records from 179 hospitals across the U.S. with 70,126 hospitalizations from 2014 to 2015— that could explain variations in clinical performance based on factors like race.

In a new paper in *PLOS Digital Health*, the investigators, including Harvineet Singh, a Ph.D. student at the NYU Center for Data Science, and Vishwali Mhasawade, a Ph.D. candidate, both under Chunara's direction, found that mortality risk prediction models that included clinical (vitals, labs and surgery) variables developed in one hospital or geographic region exhibited a lack of generalizability to different hospitals or regions. Based on a causal discovery analysis, they postulated that this lack of generalizability results from dataset shifts in race and clinical variables across hospitals or regions. In short, the race variable is intimately connected to clinical variables.

"It is clear from this research that data models—in terms of factors like mortality risk prediction at a hospital to hospital and regional hospital group level—are not immediately generalizable, and that has implications for hospitals that can't generate these models for themselves," said Chunara.

Findings also demonstrate evidence that predictive models can exhibit



disparities in performance across <u>racial groups</u> even while performing well in terms of average population-wide metrics.

"While it is well documented that clinical factors and outcomes can vary significantly by race, it is critical that we understand why those differences exist, and thus examination of data and models must be done in a larger context alongside diverse influences, from geographic and socioeconomic to clinical," she said.

Specifically, the study suggests that beyond algorithmic fairness metrics, an understanding of data generating processes for sub-groups is needed to identify and mitigate sources of variation, and to decide whether to use a risk prediction model in new environments.

**More information:** Harvineet Singh et al, Generalizability challenges of mortality risk prediction models: A retrospective analysis on a multi-center database, *PLOS Digital Health* (2022). <u>DOI:</u> <u>10.1371/journal.pdig.0000023</u>

## Provided by NYU Tandon School of Engineering

Citation: Researchers study how to unlock clinical risk-prediction models so they can be applied to multiple clinical settings (2022, April 12) retrieved 18 July 2024 from <u>https://medicalxpress.com/news/2022-04-clinical-risk-prediction-multiple.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.