


Recently discharged patients pose risk for spreading C. difficile infection

April 15 2022



Asymptomatic patients recently discharged from a hospital may be a

source of household transmission of *Clostridioides difficile* infection (CDI), according to a study published in the May issue of *Emerging Infectious Diseases*.

Aaron C. Miller, from University of Iowa in Iowa City, and colleagues evaluated whether hospitalized patients without diagnosed CDI increased the risk for CDI among their [family members](#) after [discharge](#). Analysis used U.S. insurance claims data (2001 to 2017).

The researchers found that CDI incidence among insurance enrollees exposed to a recently hospitalized family member was 73 percent greater than enrollees not exposed, and incidence increased with length of hospitalization among family members. There was a dose-response relationship observed between total days of within-household hospitalization and CDI incidence rate ratio. The incidence rate ratio increased from 1.30 for one to three days of hospitalization to 2.45 for >30 days of hospitalization, compared to persons whose family members were hospitalized less than one day.

"Asymptomatic *C. difficile* carriers discharged from hospitals could be a major source of community-associated CDI cases and should be considered by public health authorities during surveillance and intervention-based investigations when a CDI outbreak occurs," the authors write.

More information: [Abstract/Full Text](#)

© 2022 [HealthDay](#). All rights reserved.

Citation: Recently discharged patients pose risk for spreading *C. difficile* infection (2022, April 15) retrieved 9 April 2024 from <https://medicalxpress.com/news/2022-04-discharged-patients-pose-difficile-infection.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.