

## **Dog-mediated rabies remains persistently endemic in Africa**



Dogs vaccinated each year and time-varying vaccination coverage. (A) Total number of dogs vaccinated by year; (B) Time series of overall coverage at the district level from the simulations (vaccinations applied at the 1km2 scale); Areas covered by vaccination campaigns in (C) a year with relatively poor spatial coverage and (D) a year with good spatial coverage. Black dots correspond to dog vaccinations from the village-level campaigns that year allocated according

April 29 2022



to our georeferenced census of dogs in the district. Credit: *Science* (2022). DOI: 10.1126/science.abn0713

Dog-mediated rabies, a fatal disease that kills tens of thousands of people in Africa every year, remains persistently endemic in affected regions. That is, despite an overall low virus prevalence, and attempts to combat the deadly pathogen.

The findings, from an important new study led by the University of Glasgow and published in *Science*, are based on information gained from tracing the fatal disease in a population of more than 50,000 <u>dogs</u> in Tanzania from 2002 until 2016.

Dog-mediated <u>rabies</u> is transmitted through bites and scratches—either from dog to dog or dog to human—and is almost 100% fatal unless immediately treated with a post-bite course of injections that prevents the infection from taking hold. The deadly zoonotic virus is responsible for tens of thousands of deaths each year, mostly affecting children, particularly in low- and middle-income countries.

The WHO leads the collective United Against Rabies to drive progress towards zero human deaths from dog-mediated rabies by 2030 because the disease remains endemic across Africa and Asia. The persistent circulation of rabies at very low prevalence in largely unvaccinated domestic dog populations is an enduring scientific enigma.

Now an international team of researchers, led by Professor Katie Hampson at the University of Glasgow, has revealed new insights into the dynamics of this deadly disease in affected regions. The findings explain how the pathogen can persist at such low levels—less than 0.15% of the dog population is infected annually—meaning that it can spread



under the radar.

The study found that the mobility and behavior of individual rabid dogs helps to ensure that the virus can disperse and circulate at such low virus prevalence. The findings showed that a small proportion of infected dogs act as "super spreaders," traveling more than 10km to bite other animals and bringing new virus lineages to different communities, thereby seeding new outbreaks.

Prof Katie Hampson said: "The virus uses a clever strategy for low level persistence, which makes its dynamics harder to both detect and predict. But these new mechanistic insights also show why scaling up dog vaccination is so necessary."

To understand the virus dynamics, researchers tracked rabies transmissions, dog population densities, disease incidence and vaccination campaigns in a large <u>population</u> of domestic dogs in Serengeti, Tanzania for more than a decade.

According to the findings, the processes that underpin the persistence and prevalence of rabies operate on scales much smaller than those typically modeled for the disease—often down to the behavior of a few individual infected dogs

Over the 14 years, the researchers were able to pinpoint hundreds of introductions from neighboring villages that were able to sustain circulation across the region. The variability in rabid dog biting behavior means that most introductions do not take off, but some of the outbreaks that do last for years, illustrating how co-circulation of different rabies virus lineages underpins <u>virus</u> persistence.

Rabies is one of the Neglected Tropical Diseases that predominantly affects poor and vulnerable populations who live in remote rural



locations. Approximately 80% of human cases occur in rural areas. Although effective treatments exist to prevent the onset of rabies following a bite, they are not readily available or accessible to those in need.

Globally, rabies deaths are severely underreported and children between the ages of 5–14 years are the most frequent victims. Managing a rabies exposure, where the average cost of rabies post-exposure vaccination exceeds \$100, can be a catastrophic financial burden on affected families, whose average daily income may be as low as \$1 to \$2 per person.

**More information:** Rebecca Mancy et al, Rabies shows how scale of transmission can enable acute infections to persist at low prevalence, *Science* (2022). <u>DOI: 10.1126/science.abn0713</u>

Provided by University of Glasgow

Citation: Dog-mediated rabies remains persistently endemic in Africa (2022, April 29) retrieved 5 May 2024 from https://medicalxpress.com/news/2022-04-dog-mediated-rabies-persistently-endemic-africa.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.