


Genetic changes differed, increased in people with Alzheimer's disease

April 20 2022



Credit: Public Domain

Inside brain cells, errors in DNA can accumulate as we age. But in patients with Alzheimer's disease, these errors—known as somatic mutations—may build up at a faster rate. A new study by investigators from Brigham and Women's Hospital and Boston Children's Hospital found that patients with Alzheimer's disease (AD) have a greater number

of somatic mutations in their brain cells and that these mutations differed from people without Alzheimer's disease. The team's results are published in *Nature*.

"As we age, neurons are known to accumulate [somatic mutations](#). In AD neurons, however, we see more [mutations](#) and DNA alterations," said lead author Michael B. Miller, MD, Ph.D., of the Department of Pathology at the Brigham. "Our results suggest that AD neurons experience genomic damage that causes immense stress on cells and creates dysfunction among them. These findings may explain why many [brain cells](#) die during AD."

The team conducted its study using single-cell whole genome sequencing of 319 hippocampal and prefrontal cortex neurons of patients with or without AD to determine the link between the number and type of somatic mutations and AD. To better understand the genomic changes that occur in AD neurons, researchers sequenced tissue DNA and discovered a greater number of mutations termed somatic single-nucleotide variants (sSNVs) in patients with AD. Theorizing that the large number of mutations is the result of increased DNA oxidation, researchers then measured 8-Oxoguanine, an indicator of oxidative stress and DNA damage, and found that AD neurons were in fact more oxidized.

Ultimately, the discovery of accumulating DNA alterations in AD neurons provides researchers with a window into molecular and cellular events in AD pathogenesis. "Our findings suggest that the sheer number of oxidative lesions and somatic mutations we observed in AD neurons may contribute to its pathology," said Miller.

The authors acknowledge two main study limitations. First, two groups were primarily studied: [patients](#) with no neurologic disease and those with advanced AD based on the Braak staging system. In the future,

researchers are eager to study the neurons of individuals with intermediate-stage AD. Second, while single-cell, [whole-genome sequencing](#) was feasible for the preliminary studies, the authors note that there are advanced methods that allow for an in-depth analysis of each strand of DNA that should be explored in the future.

"In the future, we are eager to elucidate how the observed mutations in AD [neurons](#) cause neuronal cell death and are dedicated to aiding in the discovery of novel treatments that target these pathways," Miller said.

More information: Christopher Walsh, Somatic genomic changes in single Alzheimer's disease neurons, *Nature* (2022). [DOI: 10.1038/s41586-022-04640-1](#).
www.nature.com/articles/s41586-022-04640-1

Provided by Brigham and Women's Hospital

Citation: Genetic changes differed, increased in people with Alzheimer's disease (2022, April 20) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-04-genetic-differed-people-alzheimer-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--