

# L.A.'s injury rate from e-scooters may exceed national rate for motorcycles

April 6 2022

---



Credit: Pixabay/CC0 Public Domain

For a recent six-year period, the injury rate for riders of electric scooters in one section of Los Angeles was higher than the national rates for riders of motorcycles, bicycles and cars, and pedestrians, new UCLA

research has found.

The study, published in the peer-reviewed journal *PLOS One*, uses data from greater Los Angeles to highlight a risk that is likely to grow globally as e-scooters proliferate. According to a 2019 McKinsey report, around the world, shareable e-scooters, which can be rented on demand using a smartphone app, could in the coming years account for 1 in 10 trips shorter than 5 miles.

"There are millions of riders now using these scooters, so it's more important than ever to understand their impact on [public health](#)," said Dr. Joann Elmore, senior author of the study and a professor of medicine in the division of general internal medicine and health services research at the David Geffen School of Medicine at UCLA. "The finding that rates of injuries from e-scooters are similar to rates for motorcycle injuries is startling.

"The ease of public access to on-demand shareable scooters and safety regulations that are still in their infancy suggest that [e-scooter](#) operators, cities and [health care providers](#) will continue to see a significant number of injuries each year."

In [a study published in 2019](#), UCLA researchers reported that people injured in e-scooter accidents often sustain fractures and head trauma requiring treatment in an emergency department. That study relied on data for 249 people treated at UCLA Health's two hospitals — Ronald Reagan UCLA Medical Center and UCLA Medical Center, Santa Monica — between Sept. 1, 2017, and Aug. 31, 2018.

The new study encompassed many more patients — 1,354 injured people who were treated at 180 UCLA outpatient clinics as well as UCLA Health emergency departments and urgent care centers — and spanned a longer period — from Jan. 1, 2014 through May 14, 2020 —

enabling them to compare data from before and after the rollout of shareable e-scooters.

"Prior to the widespread introduction of shareable e-scooters in 2018, there were at most 13 e-scooter injuries per year," the study's authors write. "After [the] introduction of shareable e-scooter operators in our region, e-scooter injuries increased to 595 and 672 in 2018 and 2019, respectively."

Those injured in e-scooter accidents were not only the vehicles' riders, but also pedestrians who were hit by moving e-scooters or who tripped over parked [e-scooters](#). The researchers found that patients often were treated for injuries to the head and extremities. Five hundred thirty-three patients sustained injuries to more than one part of the body, 72 were admitted to the hospital, 21 were sent to the critical care unit and two died from their injuries.

"Overall, 33% of victims required substantial subsequent therapeutic clinical resources from our health system beyond a single clinical visit," the researchers write. "Therefore, the impact of novel e-scooter technology may have been underestimated by [early studies](#) of [\[emergency department\]](#) visits alone."

The estimated injury rate amounts to 115 injuries per 1 million e-scooter trips. By contrast, the national injury rates for other modes of transportation are 104 injuries per million motorcycle trips, 15 injuries per million bicycle trips, 8 injuries per million passenger car trips, and 2 injuries per million walking trips, according to a 2007 study in the *American Journal of Epidemiology*.

The researchers note some important limitations to their findings. For instance, they captured only data from UCLA Health facilities and did not include people who were treated at other hospitals and clinics,

indicating that injury numbers may be higher. And although the study is based on one specific geographic area, the authors write that their estimate of the number of injuries per million e-scooter trips is "of the same order" as findings from other limited regional studies.

"It is important to note that e-scooter injuries may be less severe and less fatal than motorcycle injuries, but we still think our e-scooter [injury](#) rate is an underestimate," said the study's first author, Dr. Kimon Ioannides, who led the study as a postdoctoral fellow at the UCLA National Clinician Scholars Program.

The study's co-authors are Pin-Chieh Wang, Kamran Kowsari, Vu, Dr. Noah Kojima, Dayna Clayton, Dr. Charles Liu, Dr. Tarak Trivedi and Dr. David Schriger, all of whom are affiliated with UCLA.

**More information:** E-scooter related injuries: Using natural language processing to rapidly search 36 million medical notes, *PLoS ONE* (2022).

Provided by University of California, Los Angeles

Citation: L.A.'s injury rate from e-scooters may exceed national rate for motorcycles (2022, April 6) retrieved 25 April 2024 from <https://medicalxpress.com/news/2022-04-la-injury-e-scooters-national-motorcycles.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.