

Middle ear fluid buildup is common in young children on ventilators, new study finds

April 4 2022



Credit: Pixabay/CC0 Public Domain

Babies and toddlers who need a tracheostomy—a tube surgically inserted into their windpipe to help relieve breathing problems—are at a high risk of accumulating fluid behind their eardrum when on a ventilator. That's



the conclusion of a new study, published in the *International Journal of Pediatric Otorhinolaryngology*, by UT Southwestern head and neck surgeons. This buildup of fluid, called a middle ear effusion, can put them at risk for ear infections, hearing loss, and delays in speech and language development.

"The majority of the time that these children are being cared for, the focus is on more pressing lung and heart concerns," said study leader Stephen R. Chorney, M.D., M.P.H., Assistant Professor of Otolaryngology-Head & Neck Surgery at UTSW and Pediatric Otolaryngologist at Children's Health. "But our study suggests we should also be mindful of things that might appear more trivial, like ear effusions, because they can impact communication skills and developmental milestones in a vulnerable population of kids."

Many young children who require a <u>tracheostomy</u> are born prematurely with underdeveloped lungs or narrow airways. In these cases, doctors may attach a <u>mechanical ventilator</u>—a form of life support to help a child breathe—to the tracheostomy tube.

Middle ear effusion (MEE) is a common problem for all <u>young children</u>. Nearly one in ten kids have ear tubes placed to help clear this fluid, treat infections, and reduce hearing loss. Dr. Chorney and his colleagues suspected, based on their own observations, that tracheostomy-dependent children on a ventilator might be at a higher risk of MEE.

The new study followed 94 children who received a tracheostomy before the age of two at Children's Medical Center Dallas between 2015 and 2020. On average, the children underwent tracheostomy at 5 months old and would then periodically obtain hearing tests to determine the presence of MEE. In the two years following their tracheostomy, 74% of children requiring mechanical ventilation developed at least one MEE while only 31% of those not on the ventilator developed a MEE. When



controlling for age, diagnosis of craniofacial syndrome, and result of newborn hearing test, mechanical ventilation predicted the presence of a MEE. Further, among all children with a tracheostomy, 80% of MEEs persisted for at least several months, between multiple hearing exams.

"This information allows us to have an objective reference point when communicating with parents," said Dr. Chorney. "We know that this is a common phenomenon and we might consider ear tubes in some of these children."

Since placing ear tubes requires general anesthesia, some children with a tracheostomy on ventilator support might not be good candidates for the procedure, and the new data helps clinicians weigh risks and benefits. In future studies, Dr. Chorney would like to explore the impact of MEEs on hearing and communication in this population.

"What we would hope is that if we're more vigilant with screening for and addressing MEEs in these children, positive outcomes on speech and language development can be achieved," he said. "But we need more data on that."

Other UTSW researchers who contributed to this study included Erin M. Wynings, Hussein Jaffal, Rachel St. John, and Romaine F. Johnson.

More information: Erin M. Wynings et al, Mechanical ventilation and middle ear effusions among tracheostomy-dependent children, *International Journal of Pediatric Otorhinolaryngology* (2022). DOI: 10.1016/j.ijporl.2022.111062

Provided by UT Southwestern Medical Center



Citation: Middle ear fluid buildup is common in young children on ventilators, new study finds (2022, April 4) retrieved 23 May 2024 from <u>https://medicalxpress.com/news/2022-04-middle-ear-fluid-buildup-common.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.