


Multicenter study of ultrasound LI-RADS suggests switch for HCC surveillance

April 6 2022



(A) Transverse gray-scale ultrasound image shows inability to visualize portion of liver. Examination received visualization score C. (B) Transverse gray-scale ultrasound image from examination performed 9 months later shows improved visualization of liver. Examination received visualization score A. Credit: American Roentgen Ray Society (ARRS), American Journal of Roentgenology (AJR)

According to ARRS' *American Journal of Roentgenology (AJR)*, findings from a multicenter study of ACR ultrasound LI-RADS visualization

scores on serial examinations may inform decisions to switch to alternative surveillance strategies in patients at risk for HCC with visualization score C.

Noting any decision should consider the number of previous examinations with score C, "only 42% of patients with visualization score C on surveillances examination have score C on follow-up examination," wrote corresponding author Aya Kamaya from the department of radiology at Stanford University School of Medicine in California.

Kamaya and colleagues' retrospective study included patients at risk for HCC who underwent at least two HCC surveillance [ultrasound](#) examinations between January 2017 and November 2020 at Stanford University Medical Center and two institutions in Dallas, TX: University of Texas Southwestern Medical Center and Parkland Health and Hospital System. Frequencies of score remaining unchanged after variable numbers of preceding examinations with the given score were determined.

Ultimately, the probability of repeat visualization score C on a subsequent examination increased with an increasing number of preceding examinations with score C. Frequency of score C was 42%, 67%, and 80% after 1, 2, and 3 consecutive preceding examinations with score C.

"Although not statistically significant," the authors of this *AJR* article acknowledged, "presence of severe [steatosis](#) and advanced [cirrhosis](#) had the highest odds ratios (2.88 and 2.38, respectively) for repeat score C in multivariable analysis."

More information: Thodsawit Tiyyarattanachai et al, Multicenter Study of ACR Ultrasound LI-RADS Visualization Scores on Serial

Examinations: Implications for Changes in Surveillance Strategies,
American Journal of Roentgenology (2022). [DOI:](#)
[10.2214/AJR.22.27405](https://doi.org/10.2214/AJR.22.27405)

Provided by American Roentgen Ray Society

Citation: Multicenter study of ultrasound LI-RADS suggests switch for HCC surveillance (2022, April 6) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-04-multicenter-ultrasound-li-rads-hcc-surveillance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.