

Providing naloxone in the emergency department can save lives

April 11 2022, by Anish K. Agarwal and Margaret Lowenstein



Credit: Unsplash/CC0 Public Domain

Overdose deaths continue to be a national tragedy. More than 800,000 Americans died from an overdose between 1999–2019, and the annual rate of overdoses has increased dramatically during the pandemic,

according to the latest data from the Centers for Disease Control and Prevention.

Roughly 7 in 10 current overdose deaths involve opioids, which means that many of them could be prevented with naloxone. Naloxone, also known by its brand name Narcan, acts to reverse [opioid overdoses](#) and save lives when used in time. It is easy to carry and use, and studies have demonstrated that laypeople can administer it safely and effectively to reverse overdoses.

But many of the people who are most likely to witness an overdose, including individuals who use opioids and their friends and [family members](#), may not have easy access to naloxone. Strategies are needed to increase uptake, carrying, and administration of naloxone, especially among at-risk individuals in the community who may not be engaged in routine health care or with community naloxone distribution efforts.

However, many at-risk individuals find themselves in the emergency departments (ED), either because of an overdose or other complications of substance use. For this reason, we recently examined the potential for ED visits as a critical, reachable moment to engage high-risk individuals in overdose prevention. As we reported recently in *JAMA Network Open*, we reached out to [at-risk patients](#) prescribed naloxone in the ED to understand whether they had obtained their naloxone during or after their ED visit, whether they were carrying it, and their plans to carry it in the future.

To collect data, we sent the patients a text messaging-based survey following their ED encounter. Of the 205 patients sent the survey, 41 (20%) completed it, a promising response rate for a historically difficult-to-reach population. The novel survey approach created a space for patients to provide input and feedback, a component important for future interventions to motivate behavior change.

The survey asked patients about their experiences and perceptions following the ED encounter related to accessing, using, and carrying naloxone. Findings included:

- Most of the patients did not carry naloxone prior to their ED, but over a third reported having a personal history of an overdose requiring naloxone and more than a quarter had used naloxone to reverse an overdose for another person in the past.
- Approximately half of the patients said that they were carrying naloxone after their ED visit and two-thirds planned to continue carrying.
- Of patients not carrying naloxone prior to their ED visit, 54% reported a plan to continue carrying in the future.

These findings on patient perspectives and experiences using naloxone indicate that EDs represent an opportunity to improve naloxone carrying. These early insights are key in helping quality improvement, operational, and administrative efforts to support addiction-based care and to save lives within the community.

The Biden Administration recently announced that for the first time, harm reduction services are a federal drug policy priority, and specifically highlighted EDs and hospitals as key settings for naloxone distribution. Although the plan included few specifics, our work suggests that bolstering funding to provide naloxone directly to patients in the ED may be an effective strategy to promote naloxone carrying and use among people who are highly likely to witness an overdose.

Increasing naloxone distribution should include not only EDs, but also extend into health centers, outpatient and inpatient settings, and nontraditional venues (e.g. libraries or public vending machines). Our work reinforces the need to focus on harm reduction methods across the spectrum of public health. This should include efforts at the local and

federal levels to sponsor widespread and low-cost access to naloxone, especially within high-risk communities. Similar to historical efforts to increase access to defibrillators (also called AEDs) within [public spaces](#) for [cardiac arrest](#), a synergistic approach to make [naloxone](#) easy and simple to access is paramount to preventing [overdose](#) deaths.

More information: Anish K. Agarwal et al, Assessment of Patient-Reported Naloxone Acquisition and Carrying With an Automated Text Messaging System After Emergency Department Discharge in Philadelphia, *JAMA Network Open* (2022). [DOI: 10.1001/jamanetworkopen.2022.3986](#)

Provided by University of Pennsylvania

Citation: Providing naloxone in the emergency department can save lives (2022, April 11) retrieved 24 April 2024 from <https://medicalxpress.com/news/2022-04-naloxone-emergency-department.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.