

# Hold the salt: Study reveals how reducing sodium intake can help patients with heart failure

April 3 2022

---



Credit: Pixabay/CC0 Public Domain

For the past century people with weak hearts have been told to lower their salt intake, but until now there has been little scientific evidence

behind the recommendation.

The largest randomized clinical trial to look at [sodium reduction](#) and heart failure reported results simultaneously in *The Lancet* and at the [American College of Cardiology's 71<sup>st</sup> Annual Scientific Session](#) over the weekend, and the findings were mixed.

Though reducing [salt intake](#) did not lead to fewer emergency visits, hospitalizations or deaths for patients with heart failure, the researchers did find an improvement in symptoms such as swelling, fatigue and coughing, as well as better overall quality of life.

"We can no longer put a blanket recommendation across all patients and say that limiting [sodium intake](#) is going to reduce your chances of either dying or being in hospital, but I can say comfortably that it could improve people's quality of life overall," said lead author Justin Ezekowitz, professor in the University of Alberta's Faculty of Medicine & Dentistry and co-director of the Canadian VIGOUR Centre.

The researchers followed 806 patients at 26 medical centers in Canada, the United States, Columbia, Chile, Mexico and New Zealand. All were suffering from heart failure, a condition in which the heart becomes too weak to pump blood effectively. Half of the study participants were randomly assigned to receive usual care, while the rest received nutritional counseling on how to reduce their dietary salt intake.

Patients in the nutritional counseling arm of the trial were given [dietitian-designed menu suggestions](#) using foods from their own region and were encouraged to cook at home without adding salt and to avoid high-salt ingredients. Most dietary sodium is hidden in processed foods or restaurant meals rather than being shaken at the table, Ezekowitz noted.

"The broad rule that I've learned from dietitians is that anything in a bag,

a box or a can generally has more salt in it than you would think," said Ezekowitz, who is also a cardiologist at the Mazankowski Alberta Heart Institute and director of the U of A's Cardiovascular Research Institute,

The target sodium intake was 1,500 milligrams per day—or the equivalent of about two-thirds of a teaspoon of salt—which is the [Health Canada](#) recommended limit for most Canadians whether they have heart failure or not.

Before the study, patients consumed an average of 2,217 mg per day, or just under one teaspoon. After one year of study, the usual care group consumed an average of 2,072 mg of sodium daily, while those who received nutritional guidance consumed 1,658 mg per day, a reduction of a bit less than a quarter teaspoon equivalent.

The researchers compared rates of death from any cause, cardiovascular hospitalization and cardiovascular emergency department visits in the two study groups but found no statistically significant difference.

They did find consistent improvements for the low-sodium group using three different quality of life assessment tools, as well as the [New York Heart Association heart failure classification](#), a measure of [heart failure](#) severity.

Ezekowitz said that he will continue to advise [heart failure](#) patients to cut back on [salt](#), but now he will be clearer about the expected benefits. He urges clinicians to recognize that dietary changes can be a useful intervention for some of their patients.

The team will do further research to isolate a marker in the blood of patients who benefited most from the low-sodium diet, with the aim of being able to give more targeted individual diet prescriptions in the future. The researchers will also follow up the trial patients at 24 months

and five years to determine whether further benefits are achieved over the long term.

**More information:** Justin A Ezekowitz et al, Reduction of dietary sodium to less than 100 mmol in heart failure (SODIUM-HF): an international, open-label, randomised, controlled trial, *The Lancet* (2022). [DOI: 10.1016/S0140-6736\(22\)00369-5](https://doi.org/10.1016/S0140-6736(22)00369-5)

Provided by University of Alberta

Citation: Hold the salt: Study reveals how reducing sodium intake can help patients with heart failure (2022, April 3) retrieved 10 September 2024 from <https://medicalxpress.com/news/2022-04-salt-reveals-sodium-intake-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.