

Alternative to open heart surgery just as effective for patients with common heart condition

May 17 2022



Credit: Unsplash/CC0 Public Domain

A study led by researchers at the National Institute for Health and Care Research (NIHR) Leicester Biomedical Research Centre has shown that a less invasive heart procedure for a common condition is just as effective as conventional open-heart surgery. The findings are published



in the *Journal of the American Medical Association (JAMA)* today (May 17).

The study compared Transcatheter Aortic Valve Implantation (TAVI) to conventional surgery in treating severe symptomatic aortic stenosis. In this condition there is a narrowing of the valve that blood flows through as it leaves the heart and treatment often requires surgery. It is estimated that 1.5% of UK adults over 55 and 3.5% of those over 75 have this condition.

The study is the first publicly funded study of its kind and involved every center in the UK that performs TAVI.

The first TAVI in the UK was performed in Leicester in 2007 by a multidisciplinary team at Glenfield Hospital in Leicester, led by Professor Jan Kovac. In a TAVI procedure, a new valve is threaded through a small tube inserted into an artery in the groin, upper arm or chest. It provides an alternative to open-heart surgery and is widely used across the UK.

Previous studies have shown TAVI is a safe and <u>effective treatment</u> for people who are not eligible for conventional surgery or at high operative risk. However, this study included patients who were at lower risk and suitable for either procedure. The study found that, one year after treatment, the rates of death from any cause and other important clinical outcomes such as stroke were similar between the TAVI group and the surgery group.

The study included 913 participants who were randomized to either undergo the TAVI procedure or conventional surgery. The average age of participants was 81. Those who underwent TAVI were more likely to experience vascular complications and require pacemakers, while those who underwent surgery were more likely to experience severe bleeding.



There was a greater improvement in symptoms, functional capacity and quality of life at 6 weeks in the patients who received TAVI and the quality of life advantage was still present at one year.

Professor William Toff, Professor of Cardiology in the University of Leicester's Department of Cardiovascular Sciences, led the UK TAVI study. He says that "it's great news for patients that TAVI is just as safe and effective as surgery even in patients at lower operative risk. This procedure is much less invasive than conventional approaches. People who would otherwise require open-heart surgery can have this procedure, and as it has a faster recovery time it means a shorter stay in hospital for patients."

"These are extremely encouraging results after one year. However, we shall continue to monitor the participants for at least 5 years to ensure no late differences in the groups emerge and to assess the durability of the TAVI valves. We hope to publish the 5-year results by the end of next year. In the meantime, patient-specific treatment recommendations should continue to be made by a multi-disciplinary heart team."

More information: Effect of transcatheter aortic valve implantation vs surgical aortic valve replacement on all-cause mortality in patients with aortic stenosis: a randomized clinical trial, *Journal of the American Medical Association* (2022). DOI: 10.1001/jama.2022.5776

Provided by University of Leicester

Citation: Alternative to open heart surgery just as effective for patients with common heart condition (2022, May 17) retrieved 11 July 2024 from https://medicalxpress.com/news/2022-05-alternative-heart-surgery-effective-patients.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.