

## Asymptomatic SARS-CoV-2 infections responsible for spreading of COVID-19 less than symptomatic infections

May 26 2022



Researchers find most SARS-CoV-2 infections were not persistently asymptomatic, and asymptomatic infections were less infectious than symptomatic infections. Credit: Monstera, Pexels (CC0, https://creativecommons.org/publicdomain/zero/1.0/)

Based on studies published through July 2021, most SARS-CoV-2



infections were not persistently asymptomatic, and asymptomatic infections were less infectious than symptomatic infections. These are the conclusions of an update of a systematic review and meta-analysis publishing May 26 in the open access journal *PLOS Medicine* by Diana Buitrago-Garcia of the University of Bern, Switzerland, and colleagues.

Debate about the level and risks of asymptomatic SARS-CoV-2 infections continues, with much ongoing research. Studies that assess people at just one time point can overestimate the proportion of true asymptomatic infections because those who go on to later develop symptoms are incorrectly classified as asymptomatic rather than presymptomatic. However, other studies can underestimate asymptomatic infections with research designs that are more likely to include symptomatic participants.

The new paper was an update of a living (as in, regularly updated) systematic review first published in April 2020, which includes additional, more recent studies through July 2021. 130 studies were included, with data on 28,426 people with SARS-CoV-2 across 42 countries, including 11,923 people defined as having asymptomatic infection. Because of extreme variability between included studies, the meta-analysis did not calculate a single estimate for asymptomatic infection rate, but it did estimate the inter-quartile range to be that 14–50% of infections were asymptomatic. Additionally, the researchers found that the secondary attack rate—a measure of the risk of transmission of SARS-CoV-2—was about two-thirds lower from people without symptoms than from those with symptoms (risk ratio 0.32, 95%CI 0.16–0.64).

"If both the proportion and transmissibility of asymptomatic infection are relatively low, people with asymptomatic SARS-CoV-2 infection should account for a smaller proportion of overall transmission than presymptomatic individuals," the authors say, while also pointing out



that "when SARS-CoV-2 community transmission levels are high, physical distancing measures and mask-wearing need to be sustained to prevent <u>transmission</u> from close contact with people with asymptomatic and presymptomatic infection."

Coauthor Nicola Low adds, "The true proportion of asymptomatic SARS-CoV-2 infection is still not known, and it would be misleading to rely on a single number because the 130 studies that we reviewed were so different. People with truly asymptomatic infection are, however, less infectious than those with symptomatic <u>infection</u>."

More information: *PLoS Medicine* (2022). <u>DOI:</u> <u>10.1371/journal.pmed.1003987</u>

Provided by Public Library of Science

Citation: Asymptomatic SARS-CoV-2 infections responsible for spreading of COVID-19 less than symptomatic infections (2022, May 26) retrieved 10 May 2024 from <u>https://medicalxpress.com/news/2022-05-asymptomatic-sars-cov-infections-responsible-covid-.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.