

Black adults' high cardiovascular disease risk not due to race itself

May 26 2022, by Kristin Samuelson



Credit: Unsplash/CC0 Public Domain

Findings from a new Northwestern Medicine study rebut the idea that Black individuals' higher risk of cardiovascular disease is because of biological differences.

Black adults are at significantly higher risk (1.6–2.4 times) for cardiovascular disease than white adults. The new study found these large differences can be explained by differences in [social determinants](#) of health (like education or neighborhood-level poverty), clinical factors (like blood pressure) and lifestyle behaviors (like dietary quality).

"The key take-home message is that racial differences in cardiovascular disease are not due to race itself, which is a social concept that is not related to biology," said corresponding author Dr. Nilay Shah, assistant professor of cardiology and epidemiology at Northwestern University Feinberg School of Medicine and a Northwestern Medicine physician. "Rather, these differences in cardiovascular disease can be explained by differences in social and clinical factors. Clinicians should be evaluating the social determinants that may be influencing the health of their patients.

"The data from this study starts to identify what contributes to the higher burden of heart disease experienced by Black adults, and how much each factor matters."

The study was published May 24 in *Circulation*, the flagship journal of the American Heart Association.

A breakdown of the findings:

Black women had a 2.4-times higher risk for cardiovascular disease compared with [white women](#). The study found that clinical factors, neighborhood-level factors and socioeconomic factors explained the largest components of the higher risk experienced by Black women.

Black men had a 1.6-times higher risk for cardiovascular disease compared with white men. The study found that clinical factors, socioeconomic factors and lifestyle behaviors explained the largest

components of the higher risk experienced by Black men.

"The findings of significantly higher risk in non-Hispanic Black adults compared with non-Hispanic white adults is not surprising—this is well-known," said senior author Dr. Sadiya Khan, assistant professor of cardiology and epidemiology at Feinberg and a Northwestern Medicine physician. "But it was surprising that the risk for cardiovascular disease was the same once social and clinical factors were considered over time. This finding is really important to rebut that there is an unexplained or genetic reason that Black individuals have higher risk."

The study's findings are important because they show that disparities in heart disease experienced by Black adults could be reduced by improving preventive care of heart disease risk factors and addressing social determinants, Shah said. The data provide a guide to identify strategies that may be particularly effective at reducing the persistent differences and disparities in heart disease that exist in the U.S.

"It is important to note that clinical risk factors, lifestyle and depression are not independent of socioeconomic status and neighborhood segregation," Khan said. "Future research needs to go upstream to target social determinants of cardiovascular health. Our study lays groundwork to help inform community-engaged interventions that ensure equal opportunities for all people to have access to high-quality foods, environments and health care."

The study evaluated data from about 5,100 Black and white adults who participated in the [CARDIA \(Coronary Artery Risk Development in Young Adults\) Study](#) at four locations in the U.S. (Chicago; Minneapolis, Minn.; Oakland, Calif.; and Birmingham, Ala.). The participants enrolled around 1985 and have been followed for over 30 years. The scientists evaluated the information participants provided starting from the time of their enrollment to determine the role of social

and clinical factors in the differences in [cardiovascular disease](#) experienced by Black compared with white adults over the course of 30 years of follow-up.

More information: Nilay S. Shah et al, Associations of Clinical and Social Risk Factors With Racial Differences in Premature Cardiovascular Disease, *Circulation* (2022). [DOI: 10.1161/CIRCULATIONAHA.121.058311](#)

Provided by Northwestern University

Citation: Black adults' high cardiovascular disease risk not due to race itself (2022, May 26) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-05-black-adults-high-cardiovascular-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.