


Study explores medical record linkage with goal of improving match accuracy

May 20 2022



Comparison of referential and probabilistic matches relative to HIE candidate pairs formed by 5 blocking schemes. (A) Probabilistic matches. Comparison of probabilistic matches relative to HIE pairs. The probabilistic method identified 1.2 M matches outside of the HIE pairs formed by 5 blocking schemes. (B) Referential matches. Comparison of referential matches relative to HIE pairs. The referential method identified 6.5 M matches outside of the HIE pairs formed by 5 blocking schemes. Fifteen thousand pairs were randomly sampled from the 324 million in-block pairs for match analysis. An additional 15 000 pairs were sampled from the 6.5 million out-of-block matches. (C) Out-of-block probabilistic and referential matches. Comparison of out-of-block matches for probabilistic and referential methods. A total of 1280 probabilistic matches were identified by the probabilistic method only, and referential method identified 5.3 M more out-of-block matches than probabilistic. Credit: *Journal of the American Medical Informatics Association* (2022). DOI: 10.1093/jamia/ocac068

Accurate linking of medical records from different sources, known as patient matching, plays a critical role in patient safety and quality of medical care, but has proven difficult to accomplish. A new study from the Regenstrief Institute is one of the first to evaluate commercially available matching methodologies in comparison to real-world gold standard data with the goal of identifying evidence-based opportunities for improving match accuracy for record linkage.

"Healthcare providers need securely shared access to all available data on an individual to make the best care decisions, yet patient data remains fragmented," said Regenstrief Institute Vice President for Data and Analytics and Indiana University School of Medicine Professor Shaun Grannis, M.D., M.S., who led the study. "Name variations, typographic or recording errors, missing information, as well as name, address and phone number changes and other issues make record linkage daunting."

To ensure effective data sharing, [electronic health records](#) (EHRs)—both within one office or hospital system and among different healthcare organizations—must accurately refer to a single, specific individual. Is the Robert Smith seen at one medical office the same person as the Robert Smith, Rob Smith or Bob Smith seeking care at a different facility? Are his health records comprehensively linked? To which Maria Garcia do various lab test results belong? Which Mike Miller was the one hospitalized for COVID-19 who requires long-term post-pandemic care?

Noting that there is no evidence-based "best" patient matching approach, the study authors compared the algorithms of probabilistic matching and the increasingly popular referential method of record linking to the gold standard of manually reviewed information from the Indiana Network for Patient Care (INPC), one of the nation's largest health information exchanges with 47 million patient registrations. The researchers found that referential patient matching, which employs data from

commercially available, non-healthcare sources, including credit header data and federal, state and local government person records, demonstrated greater sensitivity and accuracy than the more traditional probabilistic approach.

The authors noted, "As the United States continues to advance a national identity strategy for healthcare, a more consistent and broadly deployed approach to objectively evaluating matching algorithms is necessary to provide transparency and support healthcare organizations in adopting evidence-based best practice guidelines for patient matching algorithms. Consequently, health IT policymakers, including the ONC [Office of the National Coordinator], should explore strategies for expanding the [evidence base](#) for real-world matching system performance and encourage development of more consistent and transparent approaches to assessing and disseminating matching system performance."

"Evaluation of Real-World Referential and Probabilistic Patient Matching to Advance Patient Identification Strategy" is published online ahead of print in *JAMIA (Journal of the American Medical Informatics Association)*.

More information: Shaun J Grannis et al, Evaluation of real-world referential and probabilistic patient matching to advance patient identification strategy, *Journal of the American Medical Informatics Association* (2022). [DOI: 10.1093/jamia/ocac068](https://doi.org/10.1093/jamia/ocac068)

Provided by Regenstrief Institute

Citation: Study explores medical record linkage with goal of improving match accuracy (2022, May 20) retrieved 12 May 2024 from <https://medicalxpress.com/news/2022-05-explores-medical-linkage-goal-accuracy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.