

# The age of outbreaks: Experts warn of more animal disease threats

June 10 2022, by Daniel Lawler and Isabelle Tourne

---



Bats are a common spreader of zoonotic diseases, which experts warn could increase due to a range of human-induced upheavals to the animal world.

With the spread of monkeypox across the world coming hot on the heels of COVID-19, there are fears that increasing outbreaks of diseases that jump from animals to humans could spark another pandemic.

While such diseases—called zoonoses—have been around for millennia, they have become more common in recent decades due to deforestation, mass livestock cultivation, [climate change](#) and other human-induced upheavals of the animal world, experts say.

Other diseases to leap from animals to humans include HIV, Ebola, Zika, SARS, MERS, [bird flu](#) and the [bubonic plague](#).

The World Health Organization said on Thursday that it is still investigating the origins of COVID, but the "strongest evidence is still around zoonotic transmission".


And with more than 1,000 monkeypox cases recorded globally over the last month, the UN agency has warned there is a "real" risk the disease could become established in dozens of countries.

The WHO's emergencies director Michael Ryan said last week that "it's not just in monkeypox"—the way that humans and animals interact has become "unstable".

"The number of times that these diseases cross into humans is increasing and then our ability to amplify that disease and move it on within our communities is increasing," he said.

## Monkeypox

Zoonotic viral disease that occurs primarily in tropical rainforest areas of Central and West Africa



A member of the *Orthopoxvirus* genus, which includes smallpox

### Origins

First identified in humans in 1970 in DR Congo

Most cases reported from rainforest regions of the **Congo Basin**, particularly in the DR Congo

Considered endemic in several African nations

Mostly transmitted to people from wild animals such as rodents and primates

Person-to-person transmission possible by contact with lesions, body fluids, respiratory droplets and contaminated materials such as bedding



Not considered to be highly contagious between people

Source: ECDC/WHO/UN

### Current outbreaks

As of June 9

#### Outside endemic regions

- ▶ More than 1,000 confirmed cases reported to WHO
- ▶ Now detected in 29 countries

"The risk of monkeypox becoming established in non-endemic countries is real": WHO chief

UNAIDS said "a significant proportion" of recent monkeypox cases has been identified among gay, bisexual and other men who have sex with men

But transmission is most likely via close physical contact and could affect anyone

#### Countries with confirmed cases



#### Endemic countries

More than 1,400 cases and 66 deaths in endemic countries so far this year

### The disease

- ▶ Incubation usually 6 to 16 days
- ▶ Symptoms usually last 2 to 4 weeks

### General symptoms

- Fever • Headache • Backache
- Muscle aches • Chills • Swollen lymph nodes

### Rash



Often begins on the face

Spreads to other parts of body, including genitals

Rash forms into lesions which form a scab and eventually fall off

- ▶ Most people recover within weeks
- ▶ Up to a tenth of persons infected may die, with most deaths occurring in younger age groups
- ▶ Smallpox vaccines have been found to be about 85% effective; however access is limited as smallpox has been eradicated

AFP

Monkeypox.

Monkeypox did not recently leap over to humans—the first human case was identified in DR Congo in 1970 and it has since been confined to areas in Central and Western Africa.

Despite its name, "the latest monkeypox outbreak has nothing to do with monkeys," said Olivier Restif, epidemiologist at the University of

Cambridge.

While it was first discovered in macaques, "zoonotic transmission is most often from rodents, and outbreaks spread by person-to-person contact," he told AFP.


## **Worse yet to come?**

Around 60 percent of all known human infections are zoonotic, as are 75 percent of all new and emerging infectious diseases, according to the UN Environment Program.

Restif said the number of zoonotic pathogens and outbreaks have increased in the past few decades due to "[population growth](#), livestock growth and encroachment into wildlife habitats".

"Wild animals have drastically changed their behaviors in response to human activities, migrating from their depleted habitats," he said.





WHO has warned there is a "real" risk that monkeypox could become established in nations where it has recently spread.

"Animals with weakened immune systems hanging around near people and [domestic animals](#) is a sure way of getting more pathogen transmission."

Benjamin Roche, a specialist in zoonoses at France's Institute of Research for Development, said that deforestation has had a major effect.

"Deforestation reduces biodiversity: we lose animals that naturally regulate viruses, which allows them to spread more easily," he told AFP.

And worse may be to come, with a major study published earlier this



year warning that climate change is ramping the risk of another pandemic.

As [animals](#) flee their warming [natural habitats](#) they will meet other species for the first time—potentially infecting them with some of the 10,000 zoonotic viruses believed to be "circulating silently" among wild mammals, mostly in tropical forests, the study said.

Greg Albery, a [disease](#) ecologist at Georgetown University who co-authored the study, told AFP that "the host-pathogen network is about to change substantially".



Hidden threat: 10,000 viruses that could infect humans are 'circulating silently' among wild mammals in tropical forests.

## 'We have to be ready'

"We need improved surveillance both in urban and [wild animals](#) so that we can identify when a pathogen has jumped from one species to another—and if the receiving host is urban or in close proximity to humans, we should get particularly concerned," he said.

Eric Fevre, a specialist in infectious diseases at Britain's University of Liverpool and the International Livestock Research Institute in Kenya, said that "a whole range of new, potentially dangerous diseases could emerge—we have to be ready".

This includes "focusing the public health of populations" in remote environments and "better studying the ecology of these natural areas to understand how different species interact".

Restif said that there is "no silver bullet—our best bet is to act at all levels to reduce the risk".

"We need huge investment in frontline healthcare provision and testing capacity for deprived communities around the world, so that outbreaks can be detected, identified and controlled without delays," he said.

On Thursday, a WHO scientific advisory group released a preliminary report outlining what needs to be done when a new zoonotic pathogen emerges.

It lists a range of early investigations into how and where the pathogen jumped to humans, determining the potential risk, as well as longer-term environmental impacts.

© 2022 AFP

Citation: The age of outbreaks: Experts warn of more animal disease threats (2022, June 10) retrieved 15 August 2024 from <https://medicalxpress.com/news/2022-06-age-outbreaks-experts-animal-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.